



The effect of seed size grading on seed use efficiency and height of one-year-old container-grown Scots pine (*Pinus sylvestris* L.) seedlings

Arthur I. Novikov¹✉ and Vladan Ivetic²

¹ Voronezh State University of Forestry and Technologies named after G.F. Morozov, Timiryazeva 8, 394087 Voronezh, Russian Federation

² University of Belgrade – Faculty of Forestry, Kneza Višeslava 1, 11030 Belgrade, Serbia

✉ arthur.novikov@vglta.vrn.ru

ARTICLE INFO

Citation:

Novikov AI, Ivetic V (2018) The effect of seed size grading on seed use efficiency and height of one-year-old container-grown Scots pine (*Pinus sylvestris* L.) seedlings. *Reforesta* 6: 100-109. DOI: <https://dx.doi.org/10.21750/REFO R.6.08.61>

Editor: Jelena Aleksić, Serbia

Received: 2018-11-29

Accepted: 2018-12-19

Published: 2018-12-28



This article was presented at the International Conference “Reforestation Challenges 2018” which was held in Belgrade, Serbia, June 20-22, 2018.

Copyright: © 2018 Novikov Arthur I., Ivetic Vladan. This work is licensed under a [Creative Commons Attribution 4.0 International Public License](https://creativecommons.org/licenses/by/4.0/).



Abstract

The seed of *Pinus sylvestris* L. originating from natural forest in Voronezh region, Russian Federation, was graded in four classes based on seed diameter size, using the operational equipment of screens with round openings. The ungraded seedlot was used as control. Seedlings were produced in Voronezh containerized forest nursery, from March to September 2015, following the standard procedure for *Pinus sylvestris*. Seed from each size class, plus control, is sown in containers with 120 cm³ cells, filled with peat-substrate, grown in greenhouse for eight weeks, followed by hardening in growing area until the end of the growing season. The results shows a week positive effect of seed grading on height of one-year old container-grown seedlings, but significant improvement of seed use efficiency by reducing the number of empty cells. Based on results of this study, we can recommend grading of *Pinus sylvestris* seed on two size classes.

Keywords

Seed grading; Seed size; Seedling quality; Container-grown seedlings

Contents

| | | |
|---|---|-----|
| 1 | Introduction | 101 |
| 2 | Material and Method | 102 |
| | 2.1 Seed grading | 102 |
| | 2.2 Seedlings production | 102 |
| | 2.3 Statistical Analysis | 103 |
| 3 | Results | 103 |
| 4 | Discussion | 106 |
| | 4.1 Is there a relationship between seed size and seedling height? | 106 |
| | 4.2 Is there a relationship between seedling/container position and seedlings height? | 106 |
| | 4.3 Does seed grading on size improve seed use efficiency? | 106 |
| 5 | Conclusions | 107 |
| 6 | Acknowledgements | 107 |
| 7 | References | 107 |

1 Introduction

Improvement of forest restoration, reforestation, and afforestation success is an imperative facing the climate change and increased frequency and intensity of natural disturbances, including strong winds, heat- and drought-induced physiological stress, forest fires, pests and diseases outbreaks. The quality of forest reproductive material (FRM) has a key role in achieving this success. This role is recognized in Strategy of Forest Complex Development until 2030 (Anonymous 2018), launched recently by the Russian Federation (September 20, 2018), which emphasize the significance of FRM production and quality.

Pinus sylvestris L. is the most spread pine species in Europe, with high importance in forestry, for pulp industry and sawn timber products. In Russian Federation, *Pinus sylvestris* is one of the most important forest-forming tree species (after larch), with constant increase of area, from 113.5 million of hectares in 1988 to 122.2 million of hectares in 2010, accounting for 15.6%, of forested land area (FAO 2012).

While the provenancing and seed collection strategy are the most important management practices in maintaining the genetic diversity (Ivetić and Devetaković 2017), the first step in improving of tree seed physical quality is grading on size. The obvious practical benefit of seed grading on size, is achieving the size uniformity for applying the mechanical sowing. Further, grading on size provide more uniform seed germination, followed by the more uniform seedling density in the seedbed. There are evidences that seed size of *Pinus* species is positively correlated to the germination, seedlings survival, and growth (Burgar 1964, Dunlap and Barnett 1983, Belcher et al. 1984) mainly through providing uniform speed of germination (Barnett 2008); although the long-term effect is disputed (Mikola 1980; Sluder 1991; Castro 1999). There are also an opposite reports on week or lack of effect of seed size on seedling development of Pine species, calling into question the justification of seed grading (Dumroese and Wenny 1987; Sluder 1991; Bladé and Vallejo 2008). In addition, there is a concern that directional selection by seed grading can change the genetic constitution of the whole seedlot (Ivetić et al. 2016).

Despite controversy and conflicting reports on effect of seed size grading, this is still a standard procedure in tree seed centers', of high practical significance especially in container nurseries equipped with mechanical sowing devices. *Pinus sylvestris* seed grading on size (e.g. in Russian Federation - Novikov 2000, 2001, 2002, 2008, 2017; Sviridov et al. 2001; Gomzyakov and Novikov 2008; Gomzyakov et al. 2010; Drapalyuk and Novikov 2018; Novikov and Novikova 2018; and in Serbia – Ivetić 2013) is done by screens with openings of different size, usually separating three to four size classes, plus impurities which are larger than openings at the top, or smaller than openings at the lowest screen; followed by gravity separation for removing of empty and undeveloped seeds, and remaining impurities, and finally by pressure-vacuum separation for removing of damaged seeds. Although existing methods for seed grading are efficient at operational level, having on mind the need for enormous quantity of FRM worldwide (Haase and Davis 2017), there is a constant need for innovative approaches (e.g. Near Infrared Spectroscopy - Farhadi et al. 2015) and development and design of new devices based on biophotonic effect (Drapalyuk and Novikov 2018).

In this study, we have tried to answer the following questions, related to one-year old container-grown *Pinus sylvestris* seedlings: 1) Is there a relationship between

seed size and seedlings height? 2) Is there a relationship between seedling/container position and seedlings height? 3) Does seed grading on size improve seed use efficiency?

2 Material and Methods

2.1 Seed grading

Cones of *Pinus sylvestris* are collected from selected seed trees in natural forest, in September 2014. This seed source is located in the Pavlovsky district of the Voronezh region, Russian Federation (Latitude 50.462169; Longitude 40.096446; Altitude 83 m asl). Seed was extracted from cones and further processed (pre-cleaning, extraction, de-winging) using the standard procedures and equipment (BCC AB, Landskrona, Sweden) in Voronezh Forest Selection and Seed Center.

From the original seedlot, four random samples of 0.5 kg were extracted: one was left as the control and the remaining three were graded on size, i.e. seed diameter (Table 1) by Cleaner & Seed Sizer (Mini-Series - BCC AB), using the screens with round openings. In the course of work, some seeds stuck in sieves were observed, which leads to a decrease in the efficiency of using the device.

Table 1. Size classes and sample size of *Pinus sylvestris* seed used in the study

| Seed size group | Seed size (mm) | Sample seed quantity |
|-----------------|-----------------|----------------------|
| 1 - Extra Small | ES ≤ 2.25 | 3 × 40 (120) |
| 2 - Small | 2.25 < S ≤ 2.75 | 3 × 40 (120) |
| 3 - Medium | 2.75 < M ≤ 3.25 | 3 × 40 (120) |
| 4 - Large | L > 3.25 | 3 × 40 (120) |
| Control | bulk | 4 × 40 (160) |

2.2 Seedlings production

Seedlings for this study were produced in Voronezh containerized forest nursery, from March to September 2015, following the standard procedure (Landis et al. 1998) for *Pinus sylvestris*: mechanized peat-substrate filling and seed sowing in containers (Hiko V-120 SS – 120 cm³), followed by growing in greenhouse for eight weeks, with automatic maintenance of humidity and temperature, and hardening in growing area with automatic irrigation until the end of the growing season. From each sees size class, a total of 120 (3 container trays X 40 cells) seeds were sown, plus 160 (4 container trays X 40 cells) seeds from the control group, resulting wit total of 640 sown cells.

In order to investigate relationship between seedling/container position and seedlings height, containers sown with seeds of different size classes (plus control) are kept together, and the same arrangement with position of each container relative to the sides of the world is maintained in the greenhouse and at the growing area (Figure 1).

Height of all established seedlings was measured at the end of growing season, from root collar diameter to the top of the seedling.




Figure 1. The arrangement with position of each container relative to other containers and to the sides of the world: S – South, N – North, E – East, W – West, C – column, R – row).

2.3 Statistical Analysis

The one-way ANOVA was used to test differences between mean values of seedlings height from different seed size classes. Descriptive statistics included number of samples, mean value, standard deviation, variance, minimum value, and maximum value. Mean values were separated using Tukey’s HSD test for unequal number of samples, with a significance level of alpha = 0.05. In order to visualize height of seedlings, as well as number of empty cells, relative to their position, the colored matrices with height values are produced.

3 Results

The seed size have significant effect on one-year old *Pinus sylvestris* seedlings height (Table 2, Table 3, and Figure 2). However, this effect is not necessarily positive. The highest mean value of height (116,3 mm) is recorded for seedlings from large seed size class (>3.25 mm), but the lowest mean value of height (98,8 mm) is recorded for seedlings from small seed size class (2.25- 2.75 mm) – much lower compared to (108,6 mm) seedlings from extra small class (<2.25 mm). The mean value of seedlings height from control group is above the average for all measured seedlings.

Table 2. Descriptive statistics (N=470) of one-year old container *Pinus sylvestris* seedlings height from different seed-size classes: N – number of seedlings, SD – standard deviation, Var – variance, min – minimum value, max – maximum value). Mean values followed by the different letter are statistically different (p<0.05).

| Size Class | HT (mm) | N | SD | Var | Min | Max |
|------------|--------------------|-----|-------|-------|-----|-----|
| Control | 113,2 ^a | 74 | 17,71 | 313,6 | 40 | 140 |
| ES | 108,6 ^a | 94 | 23,12 | 534,4 | 33 | 151 |
| S | 98,8 ^b | 90 | 23,19 | 538,0 | 20 | 142 |
| M | 111,7 ^a | 106 | 24,31 | 590,9 | 36 | 160 |
| L | 116,3 ^a | 106 | 22,49 | 506,0 | 60 | 183 |
| All Groups | 109,9 | 470 | 23,20 | 538,1 | 20 | 183 |

Table 3. Analysis of Variance (One-Way ANOVA) of one-year old container *Pinus sylvestris* seedlings height from different seed-size classes.

| | SS | df | MS | SS | df | MS | F | p |
|---------|----------|----|----------|----------|-----|----------|----------|----------|
| HT (mm) | 16734,89 | 4 | 4183,722 | 235645,2 | 465 | 506,7639 | 8,255762 | 0,000002 |




Figure 2. Containers with one-year old *Pinus sylvestris* seedlings from different seed size class (left container) and control (right container at each photo): upper left – extra small (≤ 2.25 mm), upper right – small (2.25- 2.75 mm), down left – medium (2.75- 3.25 mm), and down right – large (> 3.25 mm) seed class.

There is no obvious pattern in seedling height distribution in containers with seedlings from different size classes (Figure 3). However, it is obvious that control shows significantly more empty cells (53.75 %), compared to extra small, small, medium, and large class (21.67%, 25%, 11.67% and 11.67%, respectively).




Figure 3. The matrix of one-year old *Pinus sylvestris* seedlings height (shades of red – lower than average, shades of yellow – around average, shades of green – above average) and empty cells (red color) distribution from different size classes.

4 Discussion

4.1 Is there a relationship between seed size and seedlings height?

There are evidences that seed size and mass have a positive effect on *Pinus sylvestris* seedling height (Mikola 1980; Reich 1994; Winsa and Bergsten 1994). Our results shows a similar effect, although seedlings from extra small seed size class are significantly taller than seedlings from small class. However, due to small differences in height of seedlings from different size classes, we cannot recommend grading of seed in more than two size classes. The boundary between these two classes should depend on seedlot provenance, since seed mass of *Pinus sylvestris* differs with latitude of stand origin (Reich et al. 1994). This weak effect of seed size on seedlings height can be explained, to some extent, by favorable and uniform growing conditions in the nursery, as previously have been reported that seed mass effect on *Pinus sylvestris* seedling height is stronger under harsh (i.e. direct seeding in the field) compared to optimal (i.e. sowing in nursery) conditions (Wennström et al. 2002). The mean value of seedlings height from the control group is above average, overtopping all seed size classes except seedlings from the large class. So, from aspect of seedlings height, and similar to findings of Dumroese and Wenny (1987) for *Pinus monticola* Dougl. ex D. Don and Bladé and Vallejo (2008) for *Pinus halepensis* Mill., small improvement in seedling height obtained from seed grading do not seems to justify the increased operational costs.

4.2 Is there a relationship between seedling/container position and seedlings height?

There is no obvious pattern of *Pinus sylvestris* seedlings height or of empty cells related to their position. This can be explained by uniformity of growing conditions achieved in both nursery greenhouse and growing area. Here it should be emphasized that this was a small scale experiment (13 container trays) which enable the easy achievement of optimal and uniform growing conditions. On large scale production, it can be expected that seedlings and container cell position are affecting both, seedlings height and number of empty cells at the end of the first growing season.

4.3 Does seed grading on size improve seed use efficiency?

From operational aspect, seed grading seems to be fully justified. More than half cells sown with seed from control (ungraded) seedlot ended empty after the first growing season. In the same time, the percent of empty cells sown with graded seed ranged from 11.67% (medium and large size class) to 25% (small size class). The reasons from this can be numerous, including: cleaning by removal of empty and damaged seeds and other impurities; efficiency of sowing machine, which depends to some extent on seed size uniformity; and improved seed germination parameters (rate, speed, uniformity). Seed grading improved the overall seed quality, which can result in higher survival and growth rates, as it is previously shown for *Pinus sylvestris* (Winsa and Bergsten 1994). In our study, seed grading on size did improved the *Pinus sylvestris* seed use efficiency, i.e. the seedling-to-seed ratio (Belcher et al. 1984).

5 Conclusions

The grading on size of *Pinus sylvestris* seed had a weak positive effect on one-year old container-grown seedlings. However, the grading on seed size significantly improved the seed use efficiency by reducing the number of empty cells after the first growing season. Based on results of this study, we can recommend grading of *Pinus sylvestris* seed on two size classes.

6 Acknowledgements

The authors acknowledged Head of Voronezh containerized forest nursery, for the opportunity to research in the nursery and his constructive feedback. The authors acknowledged comments and reviews from the reviewers.

7 References

- Anonymous (2018) Strategy of development of the forest complex of the Russian Federation till 2030. [In Russian]. Retrieved 01 October 2018. <http://static.government.ru/media/files/cA4eYSe0MObgNpm5hSavTdxID77KCTL.pdf>.
- Barnett JP (2008) Relating seed treatments to nursery performance: experience with southern pines. In: Dumroese RK, Riley LE, technical coordinators. National Proceedings: Forest and Conservation Nursery Associations—2007. Fort Collins (CO): USDA Forest Service, Rocky Mountain Research Station. Proceedings RMRS-P-57:27-37. Available at: http://www.fs.fed.us/rm/pubs/rmrs_p057.html
- Belcher EW, Leach GN, Gresham HH (1984) Sizing slash pine seeds as a nursery procedure. *Tree Planters' Notes* 35(2):5-10. [Google Scholar](#)
- Bladé C, Vallejo VR (2008) Seed mass effects on performance of *Pinus halepensis* Mill. seedlings sown after fire. *Forest Ecology and Management* 255: 2362–2372. <https://doi.org/10.1016/j.foreco.2007.12.039>
- Burgar RJ (1964) The effect of seed size on germination, survival, and initial growth in white spruce. *For. Chron.* 40(1): 93-97. <https://doi.org/10.5558/tfc40093-1>
- Castro J (1999) Seed mass versus seedling performance in Scots pine: A maternally dependent trait. *New Phytologist* 144(1): 153-161. <https://doi.org/10.1046/j.1469-8137.1999.00495.x>
- Drapalyuk MV, Novikov AI (2018) Operating mechanized technologies of seeds grading in artificial reforestation: foresight analysis. *Forestry Engineering Journal [Lesotekhnicheskiiy Zhurnal - in Russian]*. 8 (4): [In Russian]
- Dumroese RK, Wenny DL (1987) Sowing sized seed of western white pine in a containerized nursery. *Western Journal of Applied Forestry* 2(4): 128-130. [Google Scholar](#)
- Dunlap JR, Barnett JP (1983) Influence of seed size on germination and early development of loblolly pine (*Pinus taeda* L.) germinants. *Can J For Res* 13: 40-44. <https://doi.org/10.1139/x83-006>
- FAO (2012) The Russian Federation forest sector - Outlook study to 2030. Food and Agriculture Organization of the United Nations Rome, 2012. Retrieved 11 December 2018 from <http://www.fao.org/docrep/016/i3020e/i3020e00.pdf>
- Farhadi M, Tigabu M, Odén P (2015) Near Infrared Spectroscopy as non-destructive method for sorting viable, petrified and empty seeds of *Larix sibirica*. *Silva Fennica* 49: article id 1340. <https://doi.org/10.14214/sf.1340>
- Gomzyakov ND, Novikov AI (2008) Mechanization of preparation of forest seeds for sowing using the latest machines and equipment [in Russian - Mekhanizatsiya podgotovki lesnykh semyan k posevu s ispol'zovaniem noveyshikh mashin i oborudovaniya]. In: Actual problems of forest complex development: materials of the international scientific and technical conference. Vologda: Vologda state technical University, pp 31–35. [In Russian] [Google Scholar](#)

- Gomzyakov ND, Tomilin AN, Novikov AI (2010) New technological complex of technical means for processing of seeds of conifers [in Russian - O novom tekhnologicheskom komplekse tekhnicheskikh sredstv dlya obrabotki semyan khvoynykh porod]. In: Severgeocotech-2010 : materials of XI International youth scientific conference, Ukhta, 17-19 March 201'. Ukhta: Ukhta state technical University, pp 123–126. [In Russian] [Google Scholar](#)
- Haase D, Davis A (2017) Developing and supporting quality nursery facilities and staff are necessary to meet global forest and landscape restoration needs. *Reforesta* 4: 69-93. <https://doi.org/10.21750/REFOR.4.06.45>
- Ivetić V, Devetaković J, Nonić M, Stanković D, Šijačić-Nikolić M (2016) Genetic diversity and forest reproductive material - from seed source selection to planting. *IForest - Biogeosciences and Forestry* 9: 801-812. <https://doi.org/10.3832/ifer1577-009>
- Ivetić V (2013) Praktikum iz Semenarstva rasadničarstva i pošumljavanja. Univerzitet u Beogradu - Šumarski fakultet, Beograd. 213 p. [in Serbian] <https://doi.org/10.13140/RG.2.1.3910.8321>
- Ivetić V, Devetaković J (2017) Concerns and evidence on genetic diversity in planted forests. *Reforesta* 3: 196-207. <https://doi.org/10.21750/REFOR.3.15.39>
- Landis TD, Tinus RW, Barnett JP (1998) The container tree nursery manual. Volume 6, Seedling propagation. Agric. Handbk. 674. Washington, DC: USDA Forest Service 166 p.
- Mikola J (1980) The effect of seed size and duration of growth on the height of Scots pine (*Pinus sylvestris* L.) and Norway spruce (*Picea abies* (L.) Karst.) provenances and progenies at the nursery stage. *Silva Fennica* 14(1) article id 5069. <https://doi.org/10.14214/sf.a15010>
- Novikov AI (2000). Some technological features of the sorting devices and development trends [in Russian - Nekotorye tekhnologicheskie osobennosti sortiroval'nykh ustroystv i tendentsii ikh razvitiya]. In: Forest and youth VSAFE - 2000: proceedings of the anniversary scientific conference of young scientists dedicated to 70-th anniversary of VSAFE. Voronezh, Russian Federation, pp 53–60. [In Russian] [Google Scholar](#)
- Novikov AI (2001) About new means of forest seeds sorting in coniferous breeds [in Russian - O novykh sposobakh sortirovaniya lesnykh semyan khvoynykh porod]. In: Forests of Eurasia in the third millennium: Proceedings of the international conference of young scientists. Moscow, Russian Federation, pp 90–91. [In Russian] [Google Scholar](#)
- Novikov AI (2002) Substantiation of the technological scheme, design and parameters of the disk separator of forest seeds [in Russian - Obosnovanie tekhnologicheskoy skhemy, konstruktsii i parametrov separatora lesnykh semyan diskovogo tipa]. Dissertation, Voronezh State University of Forestry and Technologies named after G.F. Morozov, Voronezh, Russian Federation. [In Russian] [Google Scholar](#)
- Novikov AI (2008) The effect of the length of the size sections of the disc separator on the completeness of forest seeds. *News of the St. Petersburg Forestry Academy [Izvestiya Sankt-Peterburgskoy Lesotekhnicheskoy Akademii - in Russian]* 184: 57–62. [In Russian] [Google Scholar](#)
- Novikov AI (2017) Disc separators in forest seed production [in Russian - Diskovye separatory semyan v lesohozyajstvennom proizvodstve]. Voronezh: VSUFT, 159 p. Retrieved from <https://elibrary.ru/item.asp?id=30583125> [In Russian]
- Novikov AI, Novikova TP (2018) Non-destructive quality control of forest seeds in globalization: problems and prospects of output innovative products. In: 18th International Scientific Conference on Globalization and its Socio-Economic Consequences. Rajecke Teplice, Slovakia: Univ. Zilina, preprint.
- Reich PB, Oleksyn J, Tjoelker MG (1994) Seed mass effects on germination and growth of diverse European Scots pine populations. *Canadian Journal of Forest Research* 24(2): 306-320. <https://doi.org/10.1139/x94-044>
- Sluder ER (1991) Seed and seedling size grading of slash pine has little effect on long-term growth of trees. *USDA For. Serv. Tree Plant. Notes* 42(3): 23–27 [Google Scholar](#)
- Sviridov LT, Arutyunyan SS, Novikov AI, Kochegarov AV (2001) Substantiation of parameters and development of the device for cleaning and sorting of forest seeds [in Russian - Obosnovanie

- parametrov i razrabotka ustroystva dlya ochistki i sortirovaniya lesnykh semyan]. Research Report, Voronezh. [In Russian] [Google Scholar](#)
- Winsa H, Bergsten U (1994) Direct seeding of *Pinus sylvestris* using microsite preparation and invigorated seed lots of different quality: 2-year results. Canadian Journal of Forest Research. 24: 77–86. <https://doi.org/10.1139/x94-012>
- Wennström U, Bergsten U, Nilsson JE (2002) Effects of seed weight and seed type on early seedling growth of *Pinus sylvestris* under harsh and optimal conditions. Scandinavian Journal of Forest Research 17(2): 118-130. <https://doi.org/10.1080/028275802753626764>