



## Use of magnesium in bareroot pine nurseries

David B South

College of Forestry, Wildlife and Environment, Auburn University, AL

✉ [southdb@auburn.edu](mailto:southdb@auburn.edu)

### Abstract

Pines with visible magnesium (Mg) deficiencies (i.e. yellow tips on needles) occur in bareroot nurseries throughout the world. The occurrence of “yellow-tips” is rare when soil pH is above 6.5 but they have occurred on sands (pH < 6.0) with less than 25  $\mu\text{g g}^{-1}$  Mg. If yellow-tips occur in the summer, the foliar content of yellow tips is usually less than 1,000  $\mu\text{g g}^{-1}$  Mg. Some nurseries do not produce “yellow-tip” seedlings when irrigation water contains sufficient Mg. Factors favoring a deficiency include low soil pH, high calcium in irrigation water, frequent fertilization with nitrogen and potassium and applying too much gypsum. Although various Mg fertilizers are available, many nursery managers apply dolomite or potassium-magnesium sulfate before sowing seeds and a few also apply magnesium sulfate in July or August. Soil tests are used to determine when to fertilize before sowing and foliage tests determine when to apply Mg to green seedlings. Nursery managers who follow S.A. Wilde’s forest-based soil recommendations may apply magnesium sulfate to green seedlings even when seedbeds contain adequate levels of Mg. When deficiency is minor, chlorosis on needle tips usually disappears before the fall equinox and, when applied at this time, Mg fertilizers have little or no effect on height growth. This paper reviews some of the past and current uses of Mg in bareroot nurseries and highlights a need for additional research.

### Keywords

Nutrition; Foliar analysis; Soil testing; Hidden hunger; Toxicity

### Contents

1	Introduction	8
2	History	8
3	Soil test	11
4	Tissue analysis	13
5	Soils	14
5.1	Soil pH	14
5.2	Organic matter	16
5.3	Nitrogen	17
5.4	Calcium	18
5.4.1.	Gypsum	18
5.4.2.	Ca/Mg ratio	20
5.5	Potassium	20
5.6	Irrigation water	21
5.7	Mycorrhiza	21
5.8	Freeze tolerance	22
6	Magnesium removed at harvest	22

### ARTICLE INFO

#### Citation:

South DB (2022) Use of magnesium in bareroot pine nurseries. *Reforesta* 13: 7-44.

DOI: <https://dx.doi.org/10.21750/REFOR.13.02.95>

Editor: **Vladan Ivetić**

Received: **2021-12-14**

Accepted: **2022-03-14**

Published: **2022-06-30**



**Copyright:** © 2020South David B.

This work is licensed under a [Creative Commons Attribution 4.0 International Public License](https://creativecommons.org/licenses/by/4.0/).



7	Deficiency	24
7.1	Visible symptoms	25
7.2	Hidden hunger	26
8	Toxicity	27
9	Operational use	29
10	Conclusions	30
11	Acknowledgments	31
12	References	31

## 1 Introduction

Magnesium (Mg) is the fifth most commonly applied nutrient when fertilizing pine seedlings (South and Zwolinski 1996). Despite the importance of Mg for photosynthesis (Grzebisz 2015), there are only a few published trials that involved testing Mg in bareroot pine nurseries. Due to a lack of research, several myths regarding Mg have emerged. To some extent, Mg can be considered a forgotten element (Cakmak and Yazici 2010). For this reason, a literature review was undertaken to establish what is known about Mg fertilization practices in bareroot pine nurseries. This review includes literature from the 20th century since most research trials with Mg were conducted prior to 2000.

[Abbreviations: AA = ammonium acetate soil extraction. AN = ammonium nitrate. AS = ammonium sulfate. DOL = dolomitic limestone. Kainit = kainite ( $\text{KMgSO}_4\text{Cl}\cdot 3\text{H}_2\text{O}$ ). Kieserite =  $\text{MgSO}_4\cdot \text{H}_2\text{O}$ . MAP = magnesium ammonium phosphate. MS = magnesium sulfate heptahydrate;  $\text{MgSO}_4\cdot 7\text{H}_2\text{O}$  or  $\text{MgSO}_4$ . SPM = langbeinite ( $\text{K}_2\text{SO}_4\cdot 2\text{MgSO}_4$ ). B = boron. Ca = calcium. Cl = chloride. Fe = iron. K = potassium.  $\text{LSD}_{05}$  = Least significant difference,  $\alpha = 0.05$ . Mg = magnesium. Mn = manganese. N = nitrogen. P = phosphorus. pH = potential hydrogen. S = sulfur. Soil pH was measured in water. All reported correlation coefficients are significant at  $\alpha = 0.05$ .]

## 2 History

In the past, temporary or “shifting” or “flying” nurseries produced one or two crops of seedlings without applying any fertilization (Fox 1904; Schenck 1907; Anderson 1949). Typically, these small nurseries were located near reforestation sites to reduce the cost of shipping. Usually only two crops of seedlings could be harvested without adding any Mg fertilizers (Toumey 1916). In Germany, however, the second crop of seedlings was sometimes fertilized with kainit (Fox 1904).

Soil fertility in “permanent” nurseries could be maintained using purchased fertilizers or organic amendments (Ross 1929). Many managers unknowingly applied enough Mg when irrigating seedbeds (McNabb and Heidbreder-Olson 1998) so that deficiency symptoms for Mg did not occur. At some permanent nurseries, kainit might be applied before sowing or between nursery rows (Fox 1904; Schenck 1907).

During the first half of the 20<sup>th</sup> century, chemical fertilizers were rarely used in forest nurseries in the United States (Sudworth 1900; Toumey 1916; Show 1930; Wahlenberg 1930; McDaniel 1931) and when they were used, most did not contain Mg. Sometimes conifers were grown in research nurseries with soil that contained less than  $5 \mu\text{g g}^{-1}$  Mg (Flaten 1938; Argetsinger 1941). Some fertilizers supplied low rates of Mg only because kainit was included to supply K (Stewart and Hite 1903). Lime was

sometimes applied in nurseries, but the recommendation was to apply it only as an ingredient in compost or in cases where soils were clearly deficient (Toumey 1916). Kainit and wood ash were used as fertilizers to add K, but few saw any need to apply Mg. In fact, Engstrom was against the use of chemical fertilizers (Spilsbury 1949). Although more than 15 chemical fertilizers were evaluated for pine (Wahlenberg 1930; Huberman 1935), MS was not among those tested since soils typically contained sufficient levels of Mg.

Over time, soil fertility at sandy nurseries declined and chlorosis appeared on pine seedlings (Tillotson 1917; Benzian 1965). In some cases, chlorosis was an Fe deficiency symptom (i.e. high soil pH) but sometimes yellow tips of needles appeared on N-fertilized conifers growing in acid, sandy seedbeds (Ingestad 1960; Benzian 1965). In England, nursery managers refer to Mg deficiency as “hard yellows” and in other regions the deficiency is known as “yellow-tip”.

After 1930, several researchers began testing Mg in nurseries and greenhouses (Table 1). Pot studies using either sand or washed sand, proved that Mg needed to be included in research solutions (Möller 1904; Addoms 1937; Ingestad 1962) or height growth would be reduced (Figure 1). In one trial, applying a nutrient solution without Mg reduced dry mass of *Pinus palustris* seedlings by 60% (Pessin 1937). Interest in Mg fertilization increased after yellow-tip symptoms (Table 2) occurred in pine nurseries in Wisconsin (Voigt et al. 1958) and New Zealand (Will 1961, 1962).

In the 20<sup>th</sup> century, there were two schools of thought regarding how much Mg should be added to nursery soils. Wilde (1938) said “the nutrient content of productive forest soils is a satisfactory criterion for establishing soil-fertility standards in forest nurseries.” As a result, Wilde set high target levels for Mg and K based on soil sampling in virgin natural stands. This explains why he varied fertilization rates depending upon species. His Mg nursery soil targets were 75, 150, and 225  $\mu\text{g g}^{-1}$  for *Pinus banksiana*, *Pinus resinosa* and *Pinus strobus*, respectively (Wilde 1938). Once he published tentative standards for conifers in nurseries, many accepted these standards without question. As a result, the idea that pine seedbeds should contain 150  $\mu\text{g g}^{-1}$  Mg continued into the 21<sup>st</sup> century (Briggs 2008; Bueno et al. 2012).

The second school of thought was led by researchers in Germany, New Zealand and the United Kingdom who based fertility recommendations on nursery experiments. German researchers believed guessing at seedling needs with soil tests was not as useful as fertilizer tests in nurseries (Wahlenberg 1930). Those from this school established a fertility test in Scotland in 1922 using five treatments, five replications and an alpha value of 0.05 (Steven 1928). Subsequent research demonstrated that MS (applied before or after sowing) could improve needle color (Will 1961; Benzian 1967) and might increase seedling growth in a few cases (Table 1).

Table 1. The effect of magnesium sulfate (MS) fertilization on heights (mm) or mass (g) of conifers in bareroot nurseries and greenhouses (GH). Benzian (1965) applied kieserite ( $MgSO_4 \cdot H_2O$ , 17% Mg) while others applied Epsom salts ( $MgSO_4 \cdot 7H_2O$ , 9.6% Mg). Soil pH measured before fertilization and needle chlorosis relates to control seedlings.

Species	Product	Control	MS	change	pH	Chlorosis	Reference
	kg ha <sup>-1</sup>	-mm-	-mm-				
<i>Pinus radiata</i>	224	188	319	+70%	5.0	Yes	Will 1961
<i>Pinus radiata</i> – GH	312	47	57	+21%	5.6	No	Mitchell 2000
<i>Pinus radiata</i>	224	190	210	+11%	5.2	No	Will 1962
<i>Pseudotsuga menziesii</i>	1,120	43	47	+9%	5.9	No	Schaedle 1959
<i>Pinus echinata</i>	672	195	211	+8%	6	No	Auten 1945
<i>Pinus elliotii</i> – GH	224	206	212	+3%	4.7	No	Steinbeck 1962
<i>Pinus taeda</i> – GH	536	272	279	+3%	5.3	No	Edwards et al. 1990
<i>Pinus contorta</i>	211	30	31	+3%	--	--	Benzian 1965
<i>Pinus contorta</i>	211	46	48	+3%	--	--	Benzian 1965
<i>Pinus banksiana</i>	1,120	stunted	stunted	0?	4.0	Yes	Voigt et al. 1958
<i>Pseudotsuga menziesii</i>	3,360	198	???	0?	5.1	No	van den Driessche 1963
<i>Pinus resinosa</i>	211	220	???	0?	6	No	Lunt 1938
<i>Pinus contorta</i>	211	38	37	-1%	--	--	Benzian 1965
<i>Pinus taeda</i>	876	257	254	-1%	5.2	No	Wall 1994
<i>Pinus contorta</i>	211	50	49	-2%	--	--	Benzian 1965
<i>Pinus contorta</i>	211	44	43	-3%	--	--	Benzian 1965
<i>Pinus resinosa</i> – GH	672	42	41	-3%	5.2	No	Cotton 1964
<i>Pseudotsuga menziesii</i>	280	213	199	-6%	5.2	No	Schaedle 1959
<i>Pinus elliotii</i> – GH	224	223.5	203	-9%	5.8	No	Steinbeck 1962
<i>Pinus contorta</i>	359	45	40	-10%	--	--	Benzian 1965
		-g-	-g-				
<i>Pinus taeda</i> – GH	466	2.0	2.5	+25%	--	No	Berenyi et al. 1971
<i>Pinus palustris</i>	1,568	13.5	16.4	+21%	--	No	Maki and Henry 1951
<i>Pinus palustris</i>	1,568	12.7	14.8	+16%	--	No	Maki and Henry 1951
<i>Pinus taeda</i> – GH	466	3.9	4.1	+5%	--	No	Berenyi et al. 1972
<i>Pinus taeda</i> – GH	466	3.45	3.5	+1%	--	No	Berenyi et al. 1972
<i>Pinus taeda</i> – GH	466	2.55	2.37	-7%	--	No	Berenyi et al. 1971
<i>Pinus taeda</i> – GH	466	4.9	4.5	-8%	--	No	Berenyi et al. 1972




Figure 1. Height growth of pine seedlings in greenhouses is reduced about 25% (dashed line) when nutrient solutions do not contain magnesium sulfate (Hobbs 1944; Purnell 1958; Goslin 1959; Murison 1960; Sucoff 1962; Payn et al. 1995) or magnesium chloride (Van Lear and Smith 1972).

Table 2. A selected list of photographs of magnesium deficiencies in pines.

Species	Location	Photo on page	Reference
<i>Pinus</i>	Nursery	41	Baule and Fricker 1970
<i>Pinus banksiana</i>	Greenhouse	25	Swan 1970
<i>Pinus banksiana</i>	Greenhouse	152	Donald 1991
<i>Pinus echinata</i>	Greenhouse	597	Hobbs 1944
<i>Pinus nigra</i>	Field	313	Bengtson 1968
<i>Pinus nigra</i>	Field	18	Binns et al. 1980
<i>Pinus ponderosa</i>	Greenhouse	96	Murison 1960
<i>Pinus radiata</i>	Greenhouse	86	Purnell 1958
<i>Pinus radiata</i>	Nursery	44	Will 1985
<i>Pinus radiata</i>	Nursery	157	Will 1961
<i>Pinus radiata</i>	Nursery	96	Davis et al. 2015
<i>Pinus sylvestris</i>	Greenhouse	50	Goslin 1959
<i>Pinus sylvestris</i>	Field	135	Van Goor 1963
<i>Pinus sylvestris</i>	Greenhouse	51	Hackskaylo et al. 1969
<i>Pinus sylvestris</i>	Field	77	Van Goor 1970
<i>Pinus taeda</i>	Greenhouse	9	Sucoff 1961
<i>Pinus taeda</i>	Nursery	63	South 2018

### 3 Soil tests

Soils contain exchangeable and non-exchangeable Mg. In some soils, the non-exchangeable portion may contain several times more Mg than the amount reported by various soil tests (Benzian and Smith 1973; Šrámek, et al. 2012). Over time, a small portion of the non-exchangeable Mg is weathered and becomes available to crops (Salmon and Arnold 1963). Various extraction methods (Mehlich 1, Mehlich 3, AA, etc.) have been used to estimate the level of exchangeable nutrients (Alva 1993; Davey 2002; Mylavarapu et al. 2002).

Extractions of identical soil samples might produce similar Mg values when using Mehlich 1 or Mehlich 3 (Mylavarapu et al. 2002) but the AA method extracts less Mg (Alva 1993; Culman et al. 2019). For Mehlich 3 tests, levels deemed “adequate” vary by crop, but in general,  $30 \mu\text{g g}^{-1}$  Mg is “adequate” while values below  $14 \mu\text{g g}^{-1}$  Mg are “very low.” Even when using the same extraction procedure, different laboratories will report different values for the same soil sample. As a result, managers who use laboratory C (Table 3) will apply less Mg to their seedbeds than managers who send samples to laboratory A or B. In contrast, those who believe  $30 \mu\text{g g}^{-1}$  Mg is “inadequate” may waste time and money on superfluous applications of Mg.

During the 20<sup>th</sup> century, Wilde (1958) and others believed, without much evidence, that nutrient ratios were more important than absolute contents. Wilde said a nursery growing *Pinus strobus* should have a 1 to 12 ratio of available N to available Mg (Wilde 1946; p 946). Although nutrient ratios are no longer deemed necessary (Kopittke and Menzies 2007) some private soil laboratories continue to use milliequivalents to calculate K/Mg and Ca/Mg ratios.

Most managers use soil tests to determine when to apply Mg fertilizers before sowing. When the soil test result is below a pre-determined “trigger” value, then fertilizer is applied. In contrast, a “target value” is used to determine how much fertilizer to apply. Since target values for pines of 50, 75 or  $225 \mu\text{g g}^{-1}$  Mg (Table 4) were not based on science (Berenyi et al. 1971; Wall 1994), the fertilizer applications

required to meet these targets could be a waste of time and money. For example, results from a trial from one nursery (Wall 1994) suggest there is no need to adopt a target value of 50  $\mu\text{g g}^{-1}$  Mg. In fact, even 12  $\mu\text{g g}^{-1}$  Mg (Mehlich 3) can be sufficient for growth of pines at some locations (Steinbeck 1962; Berenyi et al. 1971; Munson 1982; Manikam and Srivastava 1980; Marx et al. 1984; Edwards et al. 1990). Target soil values vary from a low of 20  $\mu\text{g g}^{-1}$  to 150  $\mu\text{g g}^{-1}$  Mg (Figure 2) but most were set without establishing nursery trials (Youngberg 1952). Target values as high as 225  $\mu\text{g g}^{-1}$  Mg were based on soil Mg values in virgin stands (Wilde 1938) while a 25  $\mu\text{g g}^{-1}$  Mg target was based on greenhouse trials (Woodwell 1958).

Table 3. Examples of magnesium soil test results (Mehlich 3) using three soil samples. Two laboratories produced similar results but laboratory C extracted three times more magnesium than the other laboratories.

Sample	Laboratory		
	A $\mu\text{g g}^{-1}$	B $\mu\text{g g}^{-1}$	C $\mu\text{g g}^{-1}$
9	8	8	30
12	13	12	30
16	9	8	26




Figure 2. For pine seedbeds, various opinions exist regarding the trigger level for fertilizing with magnesium (Mg) (White et al. 1980). Some agronomists set 60  $\mu\text{g g}^{-1}$  Mg as a satisfactory level (Hardy et al. 2014) while others use a 27  $\mu\text{g g}^{-1}$  trigger value (blue arrow – line). Another group suggests no need to add Mg to pine seedbeds when soil levels exceed 24  $\mu\text{g g}^{-1}$  (black arrow – line; South and Davey 1983) while a fourth group says 52  $\mu\text{g g}^{-1}$  Mg is sufficient (green arrow - line). Some (Wilde 1958; Slone et al. 1979; White et al. 1980; van den Driessche 1984; Briggs 2008; Bueno et al. 2012) believed Mg should be applied when soil extraction levels are below 150  $\mu\text{g g}^{-1}$  Mg (red arrow - line). For pine seedlings growing in soil, published data do not show a significant growth response to Mg application when seedlings have a normal green color with no “yellow tip” needles (Table 1).

Table 4. The concentration of soil Mg recommended for growing pine seedlings varies by individual, soil texture and species. For example, Wilde (1938) recommended  $75 \mu\text{g g}^{-1}$  for *Pinus banksiana* and  $225 \mu\text{g g}^{-1}$  for *Pinus strobus* while Sadreika (1976) recommended  $18 \mu\text{g g}^{-1}$  for *Pinus banksiana* and  $60 \mu\text{g g}^{-1}$  for *Pinus strobus*. Some values were “tentative” and subject to modification (Solan et al. 1979).

Soil texture	Minimum $\mu\text{g g}^{-1}$	Recommended $\mu\text{g g}^{-1}$	Upper value $\mu\text{g g}^{-1}$	Reference
Sandy loam	-	18	-	Sadreika 1976
Sandy loam	20	30	35	Knight 1978b
Loamy sand	25	-	30	May 1984
Sand; loamy sand	25	-	-	South and Davey 1983
Sand; loamy sand	25	-	-	Aldhous and Mason 1994
Loam	30	-	45	May 1984
Sandy loam	-	36	-	Hallett 1980
Silt loam	40	-	-	South and Davey 1983
Sandy loam	-	50	-	Kormanik et al. 1994
Sand	25	50	100	Woodwell 1958
Sandy loam	-	60	-	Sadreika 1976
Sandy loam	-	75	-	Wilde 1938
Loamy sand; sandy loam	60	-	180	Stoekeler and Jones 1957
Sand; sandy loam	96	-	180	Youngberg 1984
Loam; sandy loam	120	-	240	Martian 1989
Loam; sandy loam	-	150	-	Landis 1988
Loam; sandy loam	-	150	-	Solan et al. 1979
Sandy loam	-	225	-	Wilde 1938
Sand; sandy loam	-	225	-	Youngberg 1952

#### 4 Tissue analysis

Tissue analysis is the preferred method to decide if Mg fertilization is needed (Carter et al. 2021) and pine seedlings may show yellow-tip needles when needles contain  $1,000 \mu\text{g g}^{-1}$  Mg or less (Leaf 1968; Morrison 1974). Diagnostics may be improved by separating the yellow-tip of needles from the basal green portion and submit both for analysis (e.g. Stone 1953; Hunter et al. 1986). Routine sampling during the summer (Figure 3) provides an early warning system for yellow-tip needles.

Foliage of bareroot pine seedlings averaged  $950 \mu\text{g g}^{-1}$  Mg in Australia (Flinn et al. 1980; Hopmans and Flinn 1983),  $750 \mu\text{g g}^{-1}$  Mg in New Zealand (Knight 1978b) and  $1,000$  to  $1,100 \mu\text{g g}^{-1}$  Mg in the southern United States (Boyer and South 1985; Starkey and Enebak 2012). At a nursery experiment in Fleet England, yellow-tip *Pinus contorta* seedlings had slightly more than  $600 \mu\text{g g}^{-1}$  Mg but those treated with  $6.7 \text{ kg ha}^{-1}$  of MS contained  $850 \mu\text{g g}^{-1}$  Mg (Aldhous and Atterson 1966). When soil was treated with  $40 \text{ kg ha}^{-1}$  of Mg, *Pinus contorta* foliage had  $900 \mu\text{g g}^{-1}$  Mg at the Wareham Nursery in England (Benzian and Smith 1973).

The minimum foliar Mg concentration for a pot trial with *Pinus radiata* was  $200 \mu\text{g g}^{-1}$  Mg (Will and Knight 1968) and the minimum for bareroot *Pinus taeda*, sampled in 1982 at a nursery in Arkansas, was  $300 \mu\text{g g}^{-1}$  Mg (Table 5). Deficient bareroot *Pinus radiata* seedlings had  $600 \mu\text{g g}^{-1}$  Mg in foliage (Will 1961) and bareroot *Pinus echinata* from the Ashe Nursery averaged  $435 \mu\text{g g}^{-1}$  Mg in foliage (Bryson 1980).




Figure 3. As *Pinus taeda* seedlings increase in mass, foliar concentrations of nutrients will sometimes decline from July to November (data from three fields provided for a bareroot nursery in Mississippi in 2020).

Due to "carbohydrate dilution," the Mg concentration in pine foliage declines during the growing season (Danielson 1966; Rowan and Steinbeck 1977; Wall 1994; Rowe 1996; Sung et al. 1997; Iyer et al. 2002; Dobrahner et al. 2004). In 1959, there was about a 50% decline for *Pinus elliotii* (Table 5) and in 2020, the decline from July to November was 0%, 20% and 47% for North, South and East fields, respectively (Figure 3).

## 5 Soils

Many soils in the United States are not sandy and therefore have adequate levels of Mg. For example, nurseries located on clay loam and sandy loam soils have topsoil with more than 100 µg g<sup>-1</sup> of exchangeable Mg (Youngberg 1958; McConnell and Klages 1969; South and Davey 1983; Marx et al. 1984). In contrast, sandy soils can have topsoil with less than 25 µg g<sup>-1</sup> Mg (Figure 4) and approximately 60 % of the pine plantations in the southern United States contain less than 30 µg g<sup>-1</sup> Mg (Mehlich 3) (NCSFNC 1991). Transplanted *Pinus taeda* seedlings grow well in soil with 12 µg g<sup>-1</sup> Mg (Edwards et al. 1990) and good growth occurs even when topsoil contains 6 or 7 µg g<sup>-1</sup> Mg (NCSFNC 1991). Since there is a negative correlation between Mg and sand content (South and Davey 1983; Figure 5), Mg fertilization is more likely to be needed for seedbeds with more than 75% sand.

### 5.1 Soil pH

In nurseries, soil Mg is positively related to soil pH (South and Davey 1983; South et al. 2017) partly because dolomite is used to increase soil pH and partly because Mg leaches when hydrogen ions are plentiful. Soils with pH greater than 6.0 often contain more than 100µg g<sup>-1</sup> of extractable Mg (Martian 1989; Bueno et al. 2012) and therefore have a low risk of developing a Mg deficiency (Landis 1996). In contrast, deficiencies may occur when soil pH is below 6.0 and when the soil contains less than 25 µg g<sup>-1</sup> Mg (Will 1963). At a nursery in Wisconsin, yellow-tips occurred when soil pH was 4.0 to 4.3 and this type of chlorosis was reduced by fertilization with limestone



(Stoeckeler 1949). Sometimes foliar Mg is positively related ( $r = 0.38$ ) to soil pH (Davis et al. 2007b).

Table 5. The foliar magnesium concentration of pine seedlings. Data compiled from Voigt 1955; May et al. 1962; Danielson 1966; Iyer et al. 1971; Knight 1978b; Bryson 1980; Flinn et al. 1980; Hart and Widdowson 1979; Baer 1984; van den Driessche 1984; Boyer and South 1985; Gleson 1989; Starkey and Enebak 2012; Hans 2013; and Januszek et al. 2014.

\* The  $6,230 \mu\text{g g}^{-1}$  value for June 1959 (May et al. 1962) might be due to a typographical error or residual fertilizer imbedded in epicuticular wax.

Species	Year-Month	samples	Foliar Magnesium ( $\mu\text{g g}^{-1}$ )		
			Mean	Minimum	Maximum
<i>Pinus banksiana</i>	1954?	32	850	599	1,049
	1969?	5	1,400	1,200	1,700
<i>Pinus caribaea</i>	1979-6	21	1,176	700	2,200
<i>Pinus contorta</i>	1968-10	53	1,300	1,000	1,700
<i>Pinus echinata</i>	1978-11	50	435	320	560
<i>Pinus elliotii</i>	1954-12	39	1,100	800	1,420
	1959-6	54	1,960	1,130	6,230*
	1959-9	54	1,190	880	1,900
	1959-12	54	970	780	1,140
<i>Pinus ponderosa</i>	1979-5	9	1,500	1,200	1,800
	1980-4	9	1,300	1,100	1,500
	1981-4	9	1,400	1,300	1,500
	1985-9	6	1,840	1,930	1,730
	1985-10	3	1,510	1,480	1,560
<i>Pinus radiata</i>	1975	18	750	500	1,130
	1971-1975	12	950	800	1,100
	2011-8	72	704	200	1,300
<i>Pinus resinosa</i>	1969?	7	1,400	1,100	1,700
<i>Pinus sylvestris</i>	2003-11	15	910	800	1,000
<i>Pinus taeda</i>	1955-1	162	860	670	1,150
	1956-1	216	1,050	780	1,750
	1957-1	216	1,160	670	2,150
	1965-8	4	950	840	1,080
	1966-2	4	630	570	730
	1982-12	41	1,000	300	2,300
	2009-7	19	1,400	1,100	1,700
	2009-10	19	1,100	700	1,600
	2010-2	19	1,100	600	1,500




Figure 4. Soil magnesium (Mehlich 3) from 16 southern nurseries in the United States varies from 5 to 175  $\mu\text{g g}^{-1}$ . Each nursery is represented by a mean (black dot) of 15 soil samples (i.e. 240 points on graph with many hidden points). Magnesium fertilizer was usually applied to fields in March when the soil test was  $< 25 \mu\text{g g}^{-1}$ .




Figure 5. Soil magnesium (Mehlich 1) from 37 nurseries in the United States varied from 5 to 291  $\mu\text{g g}^{-1}$  Mg (54 samples collected from 1977 to 1980). Each nursery is represented by one to four dots (1 point per mycorrhizal study). Before sowing, MS was applied to three fields and lime was applied to 14 fields (Marx et al. 1984). The nursery with the lowest sand content was at Natchez, Mississippi and the nursery at Chiefland, Florida had the most sand. The nurseries with 5  $\mu\text{g g}^{-1}$  and 291  $\mu\text{g g}^{-1}$  were at Ft. Towson, Oklahoma and Umpqua, Oregon, respectively. Sixteen fields (out of 54 points) had soil with less than 25  $\mu\text{g g}^{-1}$  Mg.

## 5.2 Organic matter

Wilde (1958, p. 364) said that soils with critically low Mg may be “safely corrected by the addition of organic remains high in bases.” For example, adding leaves (2 cm depth) at the Vallonia Nursery (Indiana) initially increased soil Mg by 73  $\mu\text{g g}^{-1}$  Mg (Davis et al. 2007a) and adding leaf mold at the Monico Nursery (Wisconsin) increased soil Mg by 50  $\mu\text{g g}^{-1}$  Mg (Wilde and Krause 1959). Although leaves may contain 2,000  $\mu\text{g g}^{-1}$  Mg, most organic amendments contain relatively low amounts of Mg. Pine bark and sawdust may contain 600 and 100  $\mu\text{g g}^{-1}$  Mg, respectively (Mexal and Fisher 1987). Therefore, even though more than 44,000  $\text{kg ha}^{-1}$  of sawdust can be applied to seedbeds, soil levels might increase by  $< 10 \mu\text{g g}^{-1}$  Mg (May 1957; Munson 1982). Likewise, incorporating peat at the FIA Nursery in Washington and at the

Toumey Nursery in Michigan did not affect soil Mg levels (Chen 1960; Koll 2009). At the Albuquerque Nursery (New Mexico), adding 43,000 kg ha<sup>-1</sup> of sawdust had no effect on foliar Mg of pine (Mexal and Fisher 1987). Although adding 500 m<sup>-3</sup> ha<sup>-1</sup> of organic material (3 parts sawdust and 1 part sewage sludge) increased soil Mg at the Wind River Nursery in Washington, it also decreased growth and survival of outplanted *Pinus ponderosa* seedlings (Coleman et al. 1987).

Some researchers test different forms of organic matter without providing a comparison of cost data. As a result, managers are often hesitant to purchase composts without knowing the potential economic benefits (Coleman et al. 1987; Rahmani et al. 2004). Since costs of transport and handling composts are high (Crowther 1950), most bareroot nursery managers do not add compost to the soil. Adding low rates of dolomitic lime is a more economical way to increase soil Mg in nurseries (Munson 1982).

### 5.3 Nitrogen

When nursery seedbeds are fertilized with ammonium fertilizers, the amount of Mg leached can increase (Wilde and Kopitke 1940; Steinbeck 1962; Deines 1973; Knight 1981; Boxman et al. 1991; Bryson and Mills 2014). When managers install unfertilized check plots, they may observe yellow-tip needles on N fertilized seedlings while slow-growing seedlings in check plots have green needles (Stoekeler and Arneman 1960). At two nurseries in Georgia, fertilization with N reduced the level of soil Mg but pine seedlings did not become Mg-deficient because soil levels in December were still greater than 20 µg g<sup>-1</sup> Mg (Figure 6).




Figure 6. Nitrogen (N) fertilization increased foliar N levels and reduced the amount of soil Mg when sampled in December (Steinbeck 1962). This produced a negative relationship between soil Mg and foliar N ( $r = -0.296$ ;  $\alpha = 0.01$ ). The lowest N treatment (112 kg ha<sup>-1</sup> of N; source was AN = ammonium nitrate) and both urea treatments (U = Uramite) were applied before sowing (April 17 - Morgan Nursery; April 12 - Page Nursery). April soil samples indicated 46 and 72 µg g<sup>-1</sup> Mg for the Morgan and Page nurseries, respectively. Both rates (224 and 448 kg ha<sup>-1</sup>) of N were applied using four top-dressings during the month of July. *Pinus elliotii* seedlings lifted from non-fertilized plots (zero N) had 1,615 µg g<sup>-1</sup> Mg in foliage at the Morgan Nursery and 1,720 µg g<sup>-1</sup> Mg at the Page Nursery. Urea fertilization resulted in “summer chlorosis” on 30 to 70 percent of seedlings from May through August. Fertilization with ammonium nitrate (AN) in July resulted in burning of foliage and increased seedling mortality (50% mortality at the Morgan Nursery and 30% at the Page Nursery). More than 70% of these seedlings remained chlorotic until cooler temperatures in October. Foliar Mg concentrations (numbers above bars) are not indicative of yellow-tip chlorosis.

At a loamy sand nursery in Wisconsin, the correlation between Mg and nitrates in the leachate was positive ( $r = 0.37$ ) (Dobrahner et al. 2004). At a nursery in New Zealand, fertilizing pines with high rates of AS increased yellow-tip symptoms but lower rates sometimes reduced chlorosis and increased foliar concentrations of Ca, K, Mg (Will 1961). At one sandy nursery in South Carolina, seedlings fertilized with N in June and July exhibited yellow-tip symptoms (0.10% Mg in foliage) while foliage on slower-growing seedlings (no N applied before August) had 0.16% Mg in foliage. Perhaps the extra N increased growth and reduced Mg levels so that yellow-tip symptoms appeared (Stoekeler and Arneman 1960). At a bareroot pine nursery with  $31 \mu\text{g g}^{-1}$  extractable Mg, fertilization with N did not reduce foliar Mg and did not induce a Mg deficiency (Wall 1994).

In greenhouse trials, fertilizing sand with solutions containing ammonium and/or nitrate increased seedling growth and reduced the Mg concentration in pine needles (Steinbeck 1962; Blackmon 1969; Malavolta et al. 1970; Swan 1970; Majid 1984). For example, fertilization with solutions containing  $100 \mu\text{g g}^{-1}$  N increased growth but higher N rates decreased growth and decreased foliar Mg (Figure 7). A similar effect was detected for soil ammonium in non-fertilized pine plantations in the southern United States. A survey of 41 sites detected a negative correlation ( $r = -0.41$ ) between soil ammonium and foliar Mg concentration for *Pinus taeda* (NCSFNC 1991).




Figure 7. Seedling shoot mass (bars and numbers), and the concentration of magnesium (Mg) in *Pinus taeda* needles (solid line) and taproots (dashed line) can be affected by the amount of urea and ammonium nitrate in nutrient solutions (Blackmon 1969). Each nitrogen treatment also contained  $50 \mu\text{g g}^{-1}$  of phosphorus and  $50 \mu\text{g g}^{-1}$  of Mg.

## 5.4 Calcium

### 5.4.1 Gypsum

Some managers notice yellow-tip needles on pine seedlings growing in fields fertilized with N and gypsum ( $\text{CaSO}_4 \bullet 2\text{H}_2\text{O}$ ). To reduce the risk of a Mg-deficiency, the rate of gypsum can be lowered (e.g.  $< 800 \text{ kg ha}^{-1}$ ) and an application of SPM can be applied before sowing. Likewise, increasing the amount of Mg in fertilizer solutions by

just 1  $\mu\text{g g}^{-1}$  could make a large difference in uptake of Mg in pine needles (Dumbroff and Michel 1967). At nurseries where gypsum treatments produced yellow-tip needles, the symptoms usually appear after mid-season and disappear before seedlings are lifted.

When gypsum ( $>4,100 \text{ kg ha}^{-1}$ ) was applied to nursery soils without any N fertilization, yellow-tips did not form on pine seedlings (Wahlenberg 1930; South 2021). When 2-0 *Pinus radiata* seedlings were already deficient in Mg, applying gypsum ( $168 \text{ kg ha}^{-1}$ ) in the spring had no effect on height growth or needle color (Will 1961). Likewise, adding gypsum to soil before sowing did not reduce growth of asymptomatic *Pinus taeda* seedlings, even with N fertilization (Table 6). Although N can lower both soil pH and soil Mg, gypsum lowers soil Mg without much effect on soil acidity (Flinn et al. 1980; Sumner et al. 1986). For example,  $280 \text{ kg ha}^{-1}$  of gypsum leaches more Mg than  $224 \text{ kg ha}^{-1}$  of AN (Figure 8).




Figure 8. Fertilization with nitrogen (N), potassium (K) and calcium (Ca) reduced the amount of magnesium in a silt loam soil at the Edwards Nursery in North Carolina (Deines 1973). The K and Ca treatments below  $54 \mu\text{g g}^{-1}$  Mg are significantly different from the K and Ca controls and N treatments below  $70 \mu\text{g g}^{-1}$  Mg are significantly different from the N control ( $\alpha=0.10$ ).

Table 6. Growth of 1-0 bareroot *Pinus taeda* seedlings (family 1-68) growing in soil amended with pine bark, calcium sulfate (gypsum) and calcium hydroxide. Soil pH, calcium (Ca) and magnesium (Mg) were from soil samples collected in November 1986 (Marx 1990). The least significant differences ( $\alpha=0.05$ ) for height, shoot dry mass and root dry mass were 3.4 cm, 3.2 g and 0.6 g, respectively. There were no differences among treatments for soil Mg (mean =  $42.5 \mu\text{g g}^{-1}$ ).

CaSO <sub>4</sub>	Ca(OH) <sub>2</sub>	pH	Soil Ca	Ca/Mg	Height	Shoot mass	Root mass
kg ha <sup>-1</sup>	kg ha <sup>-1</sup>		$\mu\text{g g}^{-1}$		cm	g	g
0	0	4.9	204	5	32.5	7.4	2.0
5,300	-	5.1	592	14	-	7.6	2.1
10,600	-	5.3	1,092	25	-	7.7	2.0
-	2,850	5.8	686	16	31.0	6.4	2.1
-	5,700	6.8	1,210	28	28.1	5.1	1.9

### 5.4.2 Ca/Mg ratio

In this review, soil nutrient ratios are determined using parts per million ( $\mu\text{g g}^{-1}$ ) and not milliequivalents. Those who prefer milliequivalents ratios may divide the Ca/Mg ratios presented here by 1.65.

Apparently, good bareroot pine seedlings were produced using soil Ca/Mg ratios of 12 (McConnell and Klages 1969; Wall 1994), 23 (Rowan 1971) and some “good” seedlings were produced with ratios > 40 (Dickson et al. 1960; Landis 1988). In contrast, stunted bareroot pine seedlings occurred in New Zealand with a Ca/Mg ratio of 4 (Will 1961). At the New Zealand nursery, chlorosis had more to do with the absolute amount of Mg ( $12 \mu\text{g g}^{-1}$  Mg) and had little to do with a hypothetical imbalance between Ca and Mg. Even so, some believe the balance between Ca and Mg is more “important” than the absolute amount of Mg in the soil [Note: use of the word “important” is subjective without a mathematical basis]. Some say once the Ca/Mg ratio in soil exceeds 10, then managers must add Mg to prevent a reduction in chlorophyll production. While this might be true at some nurseries, there are several examples where ratios were greater than 10 and green seedlings were grown without Mg fertilization (Wall 1994; Table 7).

Some have questioned the science behind a “Ca/Mg balance” for over a century (Lipman 1916) and data exist to cast doubt on the importance of a low Ca/Mg ratio. Although a soil with a Ca/Mg ratio of 11 can certainly produce yellow-tip needles, it is false logic to assume that all soils with a Ca/Mg ratio above 10 will produce Mg-deficient pine seedlings. Even Wilde questioned the Ca/Mg ratios proposed by Moser (1933). Wilde (1946; p 83) said no concrete observations verifying such an assumption have thus far been reported in relation to tree growth.

Nursery managers successfully grew pines with Ca/Mg ratios as high as 67 (Table 7). In fact, low Ca/Mg ratios in greenhouse trials showed a reduction in early root growth (Lyle and Adams 1971). Several agronomists contend the “ideal” nutrient balance theory is flawed (Schulte and Kelling 1985; Kopittke and Menzies 2007; Gaspar and Laboski 2016; Chaganti et al. 2021).

Table 7. Growth of bareroot pine seedlings growing in soil with less than  $9 \mu\text{g g}^{-1}$  magnesium (Mehlich 1). Soil pH (water), calcium (Ca), magnesium (Mg), and potassium (K) were from soil samples collected just prior to sowing. A low dose of Mg was applied over the top of *Pinus elliottii* seedlings. Shoot height (Ht), root-collar-diameter (RCD), shoot + root mass, and natural ectomycorrhizal short roots (# per 100 short roots) were measured after lifting (Munson 1982; Marx et al. 1984). Seedling mass values (Mass) are green weights except for the dry-mass value of 5.7 g highlighted in **bold**.

Species	Nursery	State	pH	Sand	Ca	Mg	K	K/Mg	Ca/Mg	Ht	RCD	Mass	Ecto
				%	$\mu\text{g g}^{-1}$	$\mu\text{g g}^{-1}$	$\mu\text{g g}^{-1}$			cm	mm	g	#
<i>P. taeda</i>	Weyerhaeuser	OK	5.2	89	89	5	51	10	18	20.3	3.9	11.4	37
<i>P. clausa</i>	Andrews	FL	4.8	94	404	6	13	2	67	20.6	2.5	6.7	24
<i>P. taeda</i>	Champion	SC	5.1	88	56	7	49	7	7	21.4	4.9	12.9	20
<i>P. taeda</i>	Weyerhaeuser	AR	5.3	87	261	7	41	6	37	23.2	4.4	13.2	33
<i>P. virginiana</i>	Vallonia	IN	4.3	82	105	8	63	8	13	20.8	4.5	21.8	19
<i>P. elliottii</i>	Archer	FL	5.5	>85	149	8	24	3	19	23.5	5.6	<b>5.7</b>	34

### 5.5 Potassium

Applying  $224 \text{ kg ha}^{-1}$  of  $\text{K}_2\text{SO}_4$  increased chlorosis of Mg deficient pine seedlings at a nursery in New Zealand (Will 1961) and high levels of exchangeable soil

K ( $> 80 \mu\text{g g}^{-1}$ ) have been correlated with Mg deficiency symptoms in pine plantations (Beets et al. 2004). When too much K fertilizers are applied, the risk of a Mg deficiency increases (Boynton and Burrell 1944; Knight 1978a; South and Davey 1983; Grzebisz 2015). At the Morgan Nursery in Georgia, five applications of KCl (each at  $112 \text{ kg ha}^{-1}$ ) reduced foliar Mg concentration in *Pinus taeda* to  $580 \mu\text{g g}^{-1}$  (Rowan 1987). Likewise, applying K before sowing lowered soil Mg at a nursery in North Carolina (Figure 8). Although yellow-tipped pine seedlings occurred when soil K/Mg ratios were 4 (Voigt et al. 1958) and 5 (Will 1961), deficiencies were not reported when K/Mg ratios before sowing were 7, 8 or 10 (Table 7).

## 5.6 Irrigation water

At some nurseries, irrigation water contains enough Mg to meet needs of seedlings (Carlson 1979; Landis 1996). Applying 60 cm of irrigation ( $2 \text{ mg L}^{-1}$  Mg) is equivalent to applying  $120 \text{ kg ha}^{-1}$  of MS (10% Mg). When pines were growing in sand ( $12 \mu\text{g g}^{-1}$  Mg) and were irrigated with water containing  $2 \text{ mg L}^{-1}$  of Mg, the concentration in needles exceeded  $2,000 \mu\text{g g}^{-1}$  Mg (Steinbeck 1962). In contrast, when pine seedlings only received rainfall, foliar Mg was less than  $1,000 \mu\text{g g}^{-1}$  (Benzian and Smith 1973). At some locations  $1,000 \text{ mm}$  of rainfall may provide only  $1\text{-}5 \text{ kg ha}^{-1}$  of Mg (Madgwick and Ovington 1959; Allen et al. 1968; Metson 1974; Hunter 1996).

At the Weyerhaeuser Nursery in Oklahoma, soil contained  $5 \mu\text{g g}^{-1}$  Mg before sowing and harvested pine seedlings averaged  $3.9 \text{ mm}$  at the root-collar (Marx et al. 1984). If the irrigation water (60 cm) contained  $5 \text{ mg L}^{-1}$  Mg, then seedlings may have been supplied with  $30 \text{ kg ha}^{-1}$  of Mg during the growing season. About 33% of irrigation water samples from southern nurseries contain more than  $4 \text{ mg L}^{-1}$  of Mg (Figure 9). At some sandy locations, irrigating non-fertilized soil for 16 years ( $632 \text{ mm yr}^{-1}$ ) might increase soil Mg by  $14 \text{ kg ha}^{-1}$  (Albaugh et al. 2014).

When growing plants in greenhouses, some believe “The calcium and magnesium ratio in the substrate solution (and in the irrigation water) should be 3 Ca to 1 Mg if expressed as meq/L or 5 Ca to 1 Mg if expressed as ppm Ca and Mg” (Bailey et al. 1999). Even so, there are insufficient data to make the Ca/Mg ratio “an evaluation factor when judging the suitability of a water for irrigation” (Ayers and Westcot 1985). For 27% of container nurseries surveyed, the Ca/Mg ratio was greater than 5 (Argo et al. 1997). To keep nursery soil “in balance,” Davey (2002) said managers must add Mg fertilizer when irrigation water contains  $>10$  times more Ca than Mg. At one *Pinus taeda* nursery, irrigation water contained  $50 \mu\text{g L}^{-1}$  Ca and  $2 \mu\text{g L}^{-1}$  Mg (McNabb and Heidbreder-Olson 1998). To avoid producing yellow-tip needles, foliar levels at that nursery were checked monthly and SPM and MS were applied over the top of seedlings in July.

## 5.7 Mycorrhiza

Typically, non-mycorrhizal pine seedlings take up Mg and do not become Mg-deficient. For example, slow-growing, non-mycorrhizal *Pinus taeda* seedlings exhibited P deficiency symptoms while needles contained  $>900 \mu\text{g g}^{-1}$  of Mg (South et al. 1988, 2018). Adequate foliar Mg concentrations were also observed for ectomycorrhizal and non-mycorrhizal seedlings in greenhouse tests (Cumming and Weinstein 1990; Walker and McLaughlin 1997; Zhang and George 2010). It is possible that activity of “rock-

eating” ectomycorrhiza can assist in converting “unavailable” Mg to ions available to pines (van Schöll et al. 2008) but there is scant evidence this occurs in nursery soils (Metson 1974; Smits and Wallander 2017). In one pine trial, foliar Mg in needles was positively related to certain types of ectomycorrhizal fungi (Leski et al. 2010).




Figure 9. The amount of Mg in irrigation water from 34 nurseries ranged from 0.06 to 18 mg L<sup>-1</sup> (McNabb and Heidbreder-Olson 1998). When the level of Mg in irrigation water is 1 mg L<sup>-1</sup>, then 60 cm of irrigation would add approximately 6 kg ha<sup>-1</sup> yr<sup>-1</sup>. At four nurseries, the Ca/Mg ratio (μg) varied from 11 to 525.

## 5.8 Freeze tolerance

After a freeze occurred at a nursery at Inchnacardoch Scotland, conifers that received no MS were “frosted” during their first winter. Seedling injury appeared to be correlated with the degree of yellowing caused by magnesium deficiency (Aldhous and Atterson 1966). Seedlings with good color and less freeze injury received 162 kg ha<sup>-1</sup> of Mg. In contrast, fertilization with MS or SPM did not increase freeze tolerance of *Pinus taeda* (Rowan 1987; Edwards et al. 1990). MS did not affect freeze tolerance of *Pinus sylvestris* growing in nutrient solutions (Christersson 1973). Likewise, MS fertilization did not seem to increase freeze tolerance of *Pseudotsuga menziesii* (Schaedle 1959).

## 6 Amount of Mg removed at harvest

Depending on species, cultural practices, and seedling age, a million pine seedlings may contain 4 to 14 kg of Mg (Knight 1978b; Flinn et al. 1980; Donald and Young 1982; South and Boyer 1983; Dobrahner et al. 2004). When 1.7 million pine seedlings ha<sup>-1</sup> are harvested, this might remove 17 kg ha<sup>-1</sup> of Mg. Likewise, harvesting a single crop of *Zea mays* grain removes about 14 kg ha<sup>-1</sup> Mg (Heckman et al. 2003). To replenish some of the harvested nutrients, some managers fertilize with more than 17 kg ha<sup>-1</sup> of Mg (Table 8).

The Mg levels in topsoil decline over time when harvest rates exceed inputs from irrigation, fertilizers and rainfall (Will and Knight 1968). Field #7, at the Ashe Nursery in Mississippi, had 74 μg g<sup>-1</sup> Mg in 1963 and 11 μg g<sup>-1</sup> in 1980 which equates to an average decline of 3.7 μg g<sup>-1</sup> yr<sup>-1</sup>. In one year, soil Mg levels decreased 18 μg g<sup>-1</sup> in Alabama (May 1957), 11 μg g<sup>-1</sup> in Georgia (Marx 1990), 25 μg g<sup>-1</sup> in Michigan (Koll



2009) and  $7 \mu\text{g g}^{-1}$  in North Carolina (Danielson 1966). At one area of the Westvaco Nursery in South Carolina, rainfall and N-fertilization of cover-crops resulted in a loss of  $30 \mu\text{g g}^{-1}$  Mg from 1983 to 1984 (Figure 10).

Table8. Examples of operational applications of magnesium sulfate (MS), sulfate potash magnesium (SPM), and dolomitic limestone (DOL) at bareroot nurseries.

Nursery	Location	Source	Mg kg ha <sup>-1</sup>	Reference
<b>Before</b>	<b>Sowing</b>			
Magnolia	Arkansas	MS	11	Marx et al. 1984
Toumey	Michigan	MS	11	Koll 2009
Orono	Canada	MS	11	Bunting 1980
Benalla	Australia	MS	18	Hopmans and Flinn 1983
Cracow	Poland	MS	20	Januszek et al. 2014
Ironhill	England	MS	34	Thomas and Jackson 1983
Coillte	Ireland	MS	37	Morrissey and O'Reilly 2002
Rotorua	New Zealand	MS	42	Knight 1981
Headley	England	MS	120	Moffat 1994
Rochester	Washington	SPM	31	Wang and Zabowski 1998
Andrews	Florida	SPM	37	Berry 1980
Archer	Florida	SPM	37	Irwin et al. 1998
Lee	Florida	SPM	37	Leach and Gresham 1983
Archer	Florida	SPM	43	Van Rees et al. 1990
Beauregard	Louisiana	SPM	49	Marx et al. 1984
Surry	Canada	SPM	49	Donald 1991
Bessey	Nebraska	SPM	50	Schmidt 1991
Andrews	Florida	SPM	62	Rodríguez-Trejo and Duryea 2003
Westvaco	South Carolina	DOL	123	Marx et al. 1984
<b>After</b>	<b>sowing</b>			
Milton	New Zealand	MS	6	Stockley 1969
SCS	Michigan	MS	2+2+2	Marx et al. 1984
Benalla	Australia	MS	5+4+4+4+5	Hopmans and Flinn 1983
Wilson	Wisconsin	MS	13+13	Dobrahner et al. 2004
Ashe	Mississippi	MS	170	Maki and Henry 1951 – research trial
Toumey	Michigan	SPM	10	Koll 2009
Nepco lake	Wisconsin	SPM	31	Marx et al. 1984
Flint River	Georgia	SPM	35	VanderSchaaf and McNabb 2004
Surry	Canada	SPM	35	Donald 1991
Andrews	Florida	SPM	25+18	Rodríguez-Trejo and Duryea 2003
Andrews	Florida	SPM	15-14-15	Berry 1980
Archer	Florida	SPM	30+30	Irwin et al. 1998
Bessey	Nebraska	SPM	25+25+25	Schmidt 1991
Kaingaroa	New Zealand	DOL	18	Will 1961 – research trial
Griffith	North Carolina	DOL	36	Hinesley and Maki 1980




Figure 10. Trends in November-extracted soil cations (Mehlich 1) at a bareroot nursery in South Carolina. Field B-1 was managed with cover-crops from 1983 to early 1989. *Pinus taeda* seed were sown in April of 1989, 1990, 1993, 1994, 1997 and 1998. Soil pH values adjacent to dots represent soil acidity in October–November. Dolomitic limestone was applied in 1988 (1,120 kg ha<sup>-1</sup>), 1991 (2,240 kg ha<sup>-1</sup>), 1992 (1,456 kg ha<sup>-1</sup>) and 1998 (1,120 kg ha<sup>-1</sup>). Magnesium sulfate and sulfate-potassium-magnesium were not applied over the top of seedlings.

## 7 Deficiency

Symptoms of Mg deficiency have occurred on fertilized pine seedlings at long-established nurseries in Australia, England, Germany, New Zealand, Sweden and the United States. In 1959, most bareroot nurseries in northern Sweden were producing seedlings with Mg deficiency symptoms (Ingestad 1960). The “yellow-tip” symptoms (e.g., Figure 11) typically occur between the summer solstice and a month after the autumn equinox (Benzian 1959). Soils prone to producing a visible deficiency include sands and loamy sands that have a low pH and low soil organic matter. Fertilization with Mg improved needle color of Mg deficient pines without increasing height or diameter growth. Likewise, when seedlings are chlorotic due to a deficiency in S (Leaf 1968; Lyle and Pearce 1968; Bolton and Benzian 1970), applying MS could improve needle color. However, when chlorosis is due to a Mn toxicity, fertilizing pine seedlings with Mg will not make needles green (Voigt et al. 1958).

Without yellow-tip symptoms and with no growth response to Mg fertilizers, a low soil test does not mean seedlings are Mg-deficient. A foliar test is more reliable than a soil extraction method that can underestimate the amount of available Mg. Even so, some authors say a soil is “deficient” when soil tests are below 150 µg g<sup>-1</sup> Mg (Table 3). Except for a few nursery trials (Will 1961; Will and Knight 1968; Wall 1994), the minimum values listed in Table 3 were not derived experimentally. In fact, when there is adequate Mg in irrigation water, good seedling growth can occur when soil contains less than 9 µg g<sup>-1</sup> Mg (Mehlich 1) (Table 6).



Figure 11. *Pinus taeda* seedlings (August 9, 2019) showing signs of Mg deficiency (foliage at  $1,000 \mu\text{g g}^{-1}$  Mg) in a bareroot nursery in South Carolina (pH = 5.7; 22 milliequivalents of cations  $\text{kg}^{-1}$ ;  $1.4 \mu\text{g g}^{-1}$  Mg in irrigation water). Soil ratios at this location were 0.6 for K/Mg and 11 for Ca/K [ $24 \mu\text{g g}^{-1}$  Mg,  $14 \mu\text{g g}^{-1}$  K and  $263 \mu\text{g g}^{-1}$  Ca, Mehlich 3]. Older needles had chlorotic tips while new needles at the shoot tip showed no chlorosis. Slow-growing seedlings (in a nearby under-fertilized check plots) had no visible symptoms of a Mg deficiency and needles contained 0.16% Mg. The chlorotic tips on the needles turned green two weeks after this photo was taken. A similar deficiency occurred at a nursery in Virginia in 2021 where yellow tips were observed on August 16, 2021 (foliage 0.07% Mg) but symptoms disappeared by August 24.

### 7.1 Visible symptoms

Chlorosis on the tips of pine needles is the clue that allows Mg deficiencies to be easily diagnosed (Leaf 1968). The chlorosis on older needle tips occurs when Mg is translocated from these needles to the newly developing needles near the top of the seedlings.

Yellow-tip symptoms appeared on conifers at two Wisconsin nurseries in 1948 (Voigt et al. 1958). According to the authors, the “stock in certain sections of the nurseries showed marked discoloration. The symptoms appeared in July or August when the outer portion of the needles exhibited a bright yellow to gold color; the portion of the needle nearest the sheath retained the normal green color. In severe cases, however, nearly the entire needle turned yellow. The yellowing was most pronounced in beds where the soil reaction was pH 4.5 or below.” The chlorotic needles of *Pinus banksiana* contained  $600 \mu\text{g g}^{-1}$  Mg and the soil (pH 4.1) contained  $7.5 \mu\text{g g}^{-1}$  Mg (Voigt et al. 1958). Green seedlings in pH 5.1 soil had  $1,800 \mu\text{g g}^{-1}$  Mg in foliage. Completely yellow seedlings were likely affected by Mn toxicity due to acid soil (< pH 4.2) and too much rain resulting in waterlogging of seedbeds (Slaton and Iyer 1974). This would explain why potted seedlings recovered when treated with

dolomite ( $1,120 \text{ kg ha}^{-1}$ ) but remained chlorotic and stunted when treated with magnesium nitrate ( $1,120 \text{ kg ha}^{-1}$ ) or  $224 \text{ kg ha}^{-1}$  of magnesium oxide (Voigt et al. 1958).

At some nurseries, yellow-tips symptoms are ephemeral and are gone by the fall equinox. For this reason, some managers were not aware of the deficiency since foliage in August was not analyzed and seedlings in November appeared normal. The reason needles recover by late September maybe due to a reduction in N and K fertilization combined with a reduction in growth rate. At one nursery in North Carolina, seedlings lifted in February had foliage with  $630 \mu\text{g g}^{-1}\text{Mg}$  (Danielson 1966) and at the Ashe Nursery (Mississippi), seedlings lifted in November in had  $435 \mu\text{g g}^{-1}\text{Mg}$  (Bryson 1980).

## 7.2 Hidden hunger

A hidden hunger exists when there are no yellow-tip symptoms but growth is increased when seedlings are fertilized with Mg. Although Mg fertilization will sometimes increase growth in pine plantations (Stone 1953; Olykan et al. 2001), examples of increased growth from MS fertilization of asymptomatic bareroot pine seedlings are rare (Table 1). Very low Mg levels will stunt growth of pine seedlings in hydroponics (Hauer-Jákli and Tränkner 2019), in containers (Will and Knight 1968) and in nurseries (Will 1961), but transitory yellow-tip symptoms may not slow growth.

To demonstrate a true hidden hunger, researchers develop growth response curves under controlled conditions (Figure 12). However, various tests conducted on asymptomatic conifer seedlings do not show a significant height growth response from fertilizing with Mg (Auten 1945; Schaedle 1959; Benzian 1965; van den Driessche 1963; Rowan 1987; Wall 1994). Perhaps soil at these nurseries contained a sufficient level of Mg (Figure 13) or perhaps irrigation supplied sufficient amounts of Mg to seedlings. It is also possible that some fertilizers contained 1 to 3% Mg and growers may have unknowingly applied 2 to  $6 \text{ kg ha}^{-1}$  of Mg to seedlings (Munson 1982; Bailey et al. 2000).

At several nurseries, pines grew well when extractable soil Mg was less than  $9 \mu\text{g g}^{-1}$  (Table 7). In greenhouse trials, fertilization with MS did not increase height growth of asymptomatic pine seedlings growing in soil with either 2.9 (Rowan 1971), 5.5 (Manikam and Sirvastava 1980) or  $15 \mu\text{g g}^{-1}$  Mg (Cotton 1964). In another trial, fertilization with Mg (source not reported) increased growth of *Pinus radiata* (Will and Knight 1968). For *Pinus palustris*, treating seedlings with  $1,568 \text{ kg ha}^{-1}$  of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  at the Ashe Nursery might have increased green mass by 16 to 21% (Maki and Henry 1951).

In general, outplanting performance is not related to the Mg content of bareroot seedlings at time of lifting (Zarger 1964; Madgwick 1975; Baer 1984; Larsen et al. 1988; van den Driessche 1991). Although there are exceptions (e.g. Will and Knight 1968), most soils at field sites have adequate Mg and uptake begins either before or soon after new roots start to grow (Baer 1984; Kelly and Barber 1991). Bareroot *Pinus echinata* needles with  $435 \mu\text{g g}^{-1}$  Mg (at planting in March) had twice that level eight months later (Bryson 1980).




Figure 12. The effect of a magnesium fertilizer (source not reported) on the growth of *Pinus banksiana* seedlings growing in sand in a greenhouse (Swan 1970). At age 16 weeks, yellow-tipped needles were noted at zero and 0.5 µg g<sup>-1</sup> treatments but chlorosis did not occur at higher solution concentrations. The hidden hunger zone may have ranged from 1 to 5 µg g<sup>-1</sup> Mg-solution for the 16-week-old seedlings.




Figure 13. The effect of magnesium (Mg) sulfate on growth and foliar Mg concentration of *Pinus taeda* seedlings at a nursery in Texas (Wall 1994). Before sowing, soil contained 31 µg g<sup>-1</sup> Mg. Beginning six weeks after sowing, applications of Epsom salts (MgSO<sub>4</sub> • 7H<sub>2</sub>O) were made every three weeks for a total of five applications. Mg treatments had no significant effect on shoot or root mass, root-collar diameter, shoot or root length (α = 0.05). Foliar Mg at lifting in January 1989 (indicated at the top of the graph) was increased by top-dressings greater than 1,000 kg ha<sup>-1</sup>.

## 8 Toxicity

In 1618, a farmer at Epsom England noticed that although his cows would not drink water from his well, the Epsom salts in the water seemed to heal scratches and rashes (Classen et al. 2004). Although little scientific evidence supports a “detoxifying effect” from water that contains Epsom salts (Deshmukh and Ray 2019), some people continue to take baths containing MS. Even so, nursery applicators should take precautions when handling MS. Since MS is a strong purgative, workers should

minimize inhalation and ingestion while preparing fertilizer solutions (Davey and McNabb 2019).

Regarding nursery stock, Wilde (1946; p 86) said “Magnesium salts in excess produce harmful effects.” It is true that too many Cl ions can kill pine seedlings (Sucoff 1962; Franklin et al. 2002; Goodrich and Jacobi 2012) but damage caused by high rates of MgCl doesn’t prove that high rates of Mg will harm pine seedlings. Greenhouse trials (Howell 1932; Sucoff 1962; Miller and Cumming 2000) and observations with nursery-grown conifers undermine the “harmful effects” assumption. Apparently, Wilde (1946) was referring to white alkaline soils (pH 8 to 9) that contain “chlorides of sodium, magnesium, and potassium, as well as calcium chloride” (Engstrom and Stoeckeler 1941; p. 45). Pines seem to be relatively tolerant of chloride-free Mg fertilizers.

Na and Cl are toxic to pine seedlings and the upper limits for irrigation water might be  $69 \mu\text{g g}^{-1}$  for Na and  $71 \mu\text{g g}^{-1}$  for Cl (Bailey et al. 1999). Even though Mg is less toxic than Na, some suggest the upper limit for Mg is less than  $69 \mu\text{g g}^{-1}$  (Landis et al. 1989; Argo et al. 1997). Although the maximum limit for Mg is not known, one nursery in California irrigated with water that contained  $113 \mu\text{g g}^{-1}$  Mg (Landis et al. 2009).

The consequences of irrigating sandy nursery soils with water containing more than  $50 \mu\text{g g}^{-1}$  Mg is not clear since most irrigation water contains less than  $40 \mu\text{g g}^{-1}$  Mg (Argo et al. 1997; Figure 9). Pine seedlings in fertilizer trials were not injured when they were fertilized using solutions that contain more than  $70 \mu\text{g g}^{-1}$  Mg (Giertych and Farrar 1961; Schomaker 1969; Brix and van den Driessche 1974). Perhaps the true concern is with a buildup of  $\text{MgCO}_3$  in soil, and not with Mg, per se. Applying too much  $\text{MgCO}_3$  will increase soil pH (Pierce et al. 1999) and too much dolomite can reduce growth of pine seedlings (Wall 1978).

A single top-dressing of MS ( $170 \text{ kg ha}^{-1}$  of Mg) did not harm *Pinus palustris* seedlings (Maki and Henry 1951) and  $210 \text{ kg ha}^{-1}$  of Mg (applied as MAP before sowing) did not harm *Pinus taeda* seedlings (Zarger 1964). Some believe it is safe to apply more than  $270 \text{ kg ha}^{-1}$  of Mg before sowing pine (White et al. 1980; Bueno 1991). Several fields at the Syracuse Nursery produce seedlings when soil contains more than  $200 \mu\text{g g}^{-1}$  Mg (Bueno 1991) and in one greenhouse study, high soil Mg ( $1,198 \mu\text{g g}^{-1}$ ) did not harm pine seedlings (Hart et al. 1980). However, adding  $1.5 \text{ kg}$  of MS  $\text{m}^{-3}$  to a bark-peat moss mix (e.g.,  $1,500 \text{ kg ha}^{-1}$  at a soil depth of 10 cm) reduced height growth of transplanted pines by 3% to 8% (Mason et al. 1995).

Although a few authors say pine needles with Mg levels greater than  $1,600 \mu\text{g g}^{-1}$  is above the “acceptable range” and “extreme” (Mellert and Göttelein 2012) this belief is not supported by the literature (Sucoff 1962; Schomaker 1969; Maxwell 1988; Landis et al. 1989). Foliar values of  $2,200 \mu\text{g g}^{-1}$  Mg were not toxic to *Pinus caribaea* (Hart et al. 1980), *Pinus elliottii* (Steinbeck 1962), *Pinus banksiana* (Swan 1970), *Pinus ponderosa* (Baer 1984), *Pinus strobus* (Iyer 1965; Schomaker 1969), *Pinus taeda* (Boyer and South 1985), *Pinus virginiana* (Sucoff 1962) or *Pinus contorta* (Goodrich and Jacobi 2012). In fact, growth of some pines was not reduced even when foliar concentrations reached  $3,700 \mu\text{g g}^{-1}$  Mg (Goodrich and Jacobi 2012). There are no data to suggest seedlings with luxury consumption of Mg are not acceptable and should be culled before outplanting.

Researchers did not report any injury when applying either SPM (Rowan 1987) or MAP to seed (Bridger et al. 1962) or seedbeds (Benzian 1967; Bean 1965;

Zarger1964; Knight 1978; Menzies et al. 2001). When applied just prior to sowing,  $MgCO_3$  might increase soil pH which could increase damping-off. For example, a pre-sowing application of  $1,120 \text{ kg ha}^{-1}$  of  $MgCO_3$  reduced seedbed densities of *Pseudotsuga* by 19 to 29% (van den Driessche 1963). However, when applied one month before sowing, even  $3,360 \text{ kg ha}^{-1}$  of  $MgCO_3$  did not affect seedling growth of *Pseudotsuga* (van den Driessche 1963). This casts doubt on the belief that high levels of  $MgCO_3$  are toxic to pines (Wilde 1946).

Although  $MgCl_2$  can certainly produce harmful effects to seedlings, this does not mean all Mg salts cause injury (Devitt et al. 2005). In fact, adding  $MgSO_4$  can increase survival of pines treated with solutions containing  $1,367 \mu\text{g g}^{-1}$  of Cl (Sucoff 1962). Instead of decreasing growth, adding  $MgSO_4$  to  $CaCl_2$  solutions increased height growth (Figure 14).

Although it is possible  $672 \text{ kg ha}^{-1}$  of MS (just prior to sowing?) reduced germination of *Pinus echinata* by 17% (Auten 1945), this non-significant ( $LSD_{05} = 47\%$ ) reduction might have been due to unrelated damping-off resulting in unusually high random variation.




Figure 14. When availability of magnesium (Mg) is low (blue line =  $0.15 \mu\text{g g}^{-1}$  Mg in solution), adding additional chloride to nutrient solutions reduces height growth of *Pinus virginiana* and too much chloride killed seedlings (Sucoff 1962). Adding additional  $MgSO_4$  (green line –  $24 \mu\text{g g}^{-1}$  Mg) reduced injury from calcium chloride. Heights are the average of four seedlings except for the container where all seedlings died (high calcium + low Mg).  $LSD_{05}$  is estimated using results from a Duncan’s multiple range test.

## 9 Costs

Fertilizer costs vary by region, shipping distance, year, and distributor. Cost comparisons, therefore, will vary over time and region. To keep Mg costs low, bareroot nursery managers in the United States typically fertilize with dolomitic limestone or SPM. When liming to increase soil pH, managers usually choose dolomitic limestone, since it is a cost-effective way to increase Mg. Assuming minimal benefit from Ca, a kg of Mg might cost \$0.30 for regular dolomite (ag grade) or \$3.00 for

pelletized dolomite. In contrast, Mg in some of the low-concentration products (Table 9) may cost more than \$100 kg<sup>-1</sup> of Mg.

At the Kaingaroa Nursery, applying a top-dressing of dolomite (168 kg ha<sup>-1</sup>) to yellow-tipped *Pinus radiata* seedlings improved seedling color (Will 1961) but the chlorosis was eliminated in plots treated with, more soluble, MS. When dolomitic lime is incorporated about a year before sowing pines, a rate of 1,000 kg ha<sup>-1</sup> will initially add about 110 kg ha<sup>-1</sup> of Mg.

Table 9. A partial list of magnesium (Mg) fertilizers. soluble (So): granules (G): liquids (L).

Ingredients	% Mg	Tradename	Form	% Mg	% S	% N
magnesium oxide + magnesium sulfate	50-55	ProMag®	G	36	6	
magnesium oxysulfate	36	Magnesium 36%	G	36	6	
magnesium sulfate	18	Southern Ag®	So	15	16	
magnesium ammonium phosphate	15	MagAmp	G	15		7
magnesium – chelate glycine	3-12	BioMin®	So	12		5.5
dolomitic limestone	8-20	Pro-Select	G	11.3	0.3	
kieserite MgSO <sub>4</sub> ·H <sub>2</sub> O	17	Amgrow-kieserite	G	15	16	
potassium magnesium sulfate	12	K-Mag® 0-0-22	G	10-11	21-22	
magnesium nitrate hexahydrate	10	Magnisal™	So	9.6		11
magnesium sulfate MgSO <sub>4</sub> ·7H <sub>2</sub> O	10	Hi-yield®	So	9.5	12.5	
magnesium sulfate MgSO <sub>4</sub> ·7H <sub>2</sub> O	10	Brant®	So	9.5	12.9	
magnesium – EDTA chelate	3-12	Axilo™	So	6		
magnesium chloride	25	NutriMag™	L	5.5		5
magnesium nitrate	10	Manni-Plex™ Mg	L	5		7
magnesium thiosulfate	24	MagThio®	L	4	10	
magnesium sulfate	10	BioMin®	L	3	3.9	1
magnesium acetate + KMg thiosulfate	3	Nutra -Boost®	L	3	6	
kainit KMgSO <sub>4</sub> Cl·3H <sub>2</sub> O	13	Magnesia-Kainit®	G	2	3	
K, S, Mg, Fe, B, Mn, Zn, Ca, P	1-3	Wood ash	-	2	0.4	
K, S, Mg, Fe, B, Mn, Zn	1	MaxiGreenII®	L	1	2	

SPM has been applied at 300 kg ha<sup>-1</sup> before sowing or four months after sowing. Assuming other nutrients in SPM have no value, the cost kg<sup>-1</sup> for Mg might be \$4.50 for SPM and \$6.00 for MS. The total amount of Mg applied to a 1-0 crop of pine seedlings can vary from 10 to 40 kg ha<sup>-1</sup> (Table 8). For pine plantations, some managers might apply 28 kg ha of Mg when foliar Mg drops below 600 to 800 µg g<sup>-1</sup> (Jokela et al. 2004).

When seedlings are green, many managers do not apply MS fertilizers (van den Driessche 1984; Donald 1991). If yellow-tip symptoms occur in the summer, treatments like MS or Mg nitrate may be applied over the top of pine seedlings. In some cases, the color improves within 10 days of treatment.

## 10 Conclusions

Due to faulty logic and a lack of Mg research, several myths have spread throughout the nursery literature. Likewise, several of the conclusions listed below should be tested using trials in bareroot seedbeds.

(1) When irrigation provides more than 18 kg ha<sup>-1</sup> of Mg, there is no need to fertilize 1-0 bareroot pine seedlings with extra Mg. [Note: When irrigation water



contains  $3\mu\text{g g}^{-1}$  Mg, then 600 mm of irrigation will supply  $18\text{ kg ha}^{-1}$  of Mg].

(2) When there is less than  $3\mu\text{g g}^{-1}$  Mg in irrigation water, applying too much N or too much gypsum can induce a Mg deficiency in pine seedlings.

(3) At most bareroot nurseries, there is no need to apply MS before sowing pines since soluble Mg will leach with rainfall and irrigation. When needed, longer-lasting Mg sources can be applied before sowing seed.

(4) The belief that  $51\mu\text{g g}^{-1}$  Mg in irrigation water will harm pine seedlings growing in sandy soil is not based on science. However,  $177\mu\text{g g}^{-1}$  of  $\text{MgCO}_3$  in irrigation water can increase soil pH, which will likely reduce growth of pines.

(5) Use of foliar tests (e.g. in August) might reduce routine use of MS at some bareroot nurseries and might increase use at other nurseries. Without foliar tests, some bareroot pine seedlings have been planted with foliage that contained less than  $7\mu\text{g g}^{-1}$  Mg.

(6) There are no data to suggest that 1-0 bareroot pine seedlings need to be fertilized with more than  $20\text{ kg ha}^{-1}$  of Mg.

(7) If the soil Ca/Mg ratio is greater than 10 and irrigation water contains less than  $2\mu\text{g g}^{-1}$  Mg, then pine seedlings might develop yellow-tip symptoms in the summer. These symptoms may disappear when N fertilization ceases and root growth increases (after the autumnal equinox).

(8) The belief that  $1,600\mu\text{g g}^{-1}$  Mg in pine needles is outside the “adequate range” (and is harmful to bareroot pine seedlings) is an opinion that is based on faulty logic.

(9) When a foliar spray of magnesium chloride turns “yellow-tip” seedlings green, the seedlings were Mg deficient. Likewise, when deficient seedlings turn green after treatment with sulfate ( $\text{H}_2\text{SO}_4$ ), the seedlings were S deficient. When a foliar spray of MS turns seedlings green, then a foliar analysis would help determine which nutrient (Mg or S) was deficient.

## 11 Acknowledgments

I thank members of the Southern Forest Nursery Management Cooperative for providing soil and foliage data from pine nurseries. I especially thank Chase Weatherly for discussions about gypsum and Mg fertilization in nursery seedbeds. I also thank Donald Cotton, Margot Wall, Jim Deines and Hoy Bryson for providing detailed Mg data in their theses. I thank Barbara Hawkins, John Mexal, and J.B. Jett for feedback on earlier drafts.

## 12 References

- Addoms RM (1937) Nutritional studies on loblolly pine. *Plant Physiology* 12(1): 199-205. <https://doi.org/10.1104/pp.12.1.199>
- Albaugh TJ, Kiser LC, Fox TR, Allen HL, Rubilar RA, Stape JL (2014) Ecosystem nutrient retention after fertilization of *Pinus taeda*. *Forest Sci* 60(6): 1131-1139. <http://dx.doi.org/10.5849/forsci.13-159>
- Aldhous JR, Atterson J (1966) Nursery investigations. *Rep For Res* 1965: 21-25. [https://www.forestresearch.gov.uk/documents/7129/FCFR\\_1965-1966.pdf](https://www.forestresearch.gov.uk/documents/7129/FCFR_1965-1966.pdf)
- Aldhous JR, Mason WL (1994) *Forest Nursery Practice*. Forestry Commission Bull. 111, London, UK. <https://www.forestresearch.gov.uk/documents/6578/FCBU111.pdf>
- Allen SE, Carlisle A, White EJ, Evans CC (1968) The plant nutrient content of rainwater. *The Journal of Ecology* 56(2): 497-504. <https://doi.org/10.2307/2258247>

- Alva AK (1993) Comparison of Mehlich 3, Mehlich 1, ammonium bicarbonate-DTPA, 1.0 M ammonium acetate, and 0.2 M ammonium chloride for extraction of calcium, magnesium, phosphorus, and potassium for a wide range of soils. *Commun Soil Sci Plan* 24(7-8): 603-612.  
<https://doi.org/10.1080/00103629309368826>
- Anderson ML (1949) Some observations on Belgian forestry. *Empire Forestry Review* 28(2): 117-130.  
<https://www.jstor.org/stable/42602429>
- Arnold MA, Struve DK (1993) Root distribution and mineral uptake of coarse-rooted trees grown in cupric hydroxide-treated containers. *HortSci* 28(10): 988-992.  
<https://doi.org/10.21273/HORTSCI.28.10.988>
- Argetsinger LM (1941) Chemical fertilizer treatments of Norway pine transplants in University of Michigan nursery. MF thesis. University of Michigan, Ann Arbor. 64  
p.<https://deepblue.lib.umich.edu/bitstream/handle/2027.42/113806/39015003283861.pdf?sequence=1>
- Argo WR, Biernbaum JA, Warncke DD (1997) Geographical characterization of greenhouse irrigation water. *HortTechnology* 7(1): 49-55. <https://doi.org/10.21273/HORTTECH.7.1.49>
- Auten JT (1945) Response of shortleaf and pitch pines to soil amendments and fertilizers in newly established nurseries in the central states. *J Agric Res* 70(12): 405-426.  
<https://babel.hathitrust.org/cgi/pt?id=uva.x004385460&view=1up&seq=6>
- Ayers RS, Westcot DW (1985) Water quality for agriculture. Irrigation and drainage paper 29. Food and Agriculture Organization of the United Nations: Rome. 174 p.  
<https://www.fao.org/3/t0234e/t0234e.pdf>
- Baer NW (1984) Nutrient content in ponderosa pine foliage: seasonal variation. *Agricultural Experiment Station Technical Bulletins*. 77. South Dakota State University. 10 p.  
[http://openprairie.sdstate.edu/agexperimentsta\\_tlb/77](http://openprairie.sdstate.edu/agexperimentsta_tlb/77)
- Bailey D, Bilderback T, Bir D (1999) Water considerations for container production of plants. NC State University Department of Horticultural Science Horticulture Information Leaflet 557: 11p.  
<https://www.clamerinforma.it/EnglishFiles/PDF/HIL557.pdf>
- Bailey DA, Nelson PV, Fonteno WC (2000) Substrates pH and water quality. NC State University Department of Horticultural Science Horticulture. Raleigh, NC: 5  
p.<http://www.nurserycropscience.info/water/source-water-quality/other-references/substrate-ph-and-water-quality.pdf>
- Baule H, Fricker C (1970) The fertilizer treatment of forest trees. BLV München.
- Bean D (1964) Artificial lighting fails to stimulate height growth of white pine seedlings in nursery studies. *Tree Planters' Notes* 64: 23-26. [https://rngr.net/publications/tpn/15-1/PDF.2004-08-13.4449/at\\_download/file](https://rngr.net/publications/tpn/15-1/PDF.2004-08-13.4449/at_download/file)
- Beets PN, Oliver GR, Kimberley MO, Pearce SH, Rodgers B (2004) Genetic and soil factors associated with variation in visual magnesium deficiency symptoms in *Pinus radiata*. *Forest EcolManag* 189(1-3): 263-279. <https://doi.org/10.1016/j.foreco.2003.08.013>
- Bengtson GW (ed) (1968) *Forest Fertilization—Theory and Practice*. Tennessee Valley Authority, Muscle Shoals, AL: 316 p. <https://catalog.hathitrust.org/Record/001516717>
- Benzian B (1959) Nutrition problems in forest nurseries. *Science of Food and Agriculture* 10(12): 637-644. <https://doi.org/10.1002/jsfa.2740101201>
- Benzian B (1965) Experiments on nutrition problems in forest nurseries. *Forestry Commission Bull* 37. p, 251. [https://www.forestresearch.gov.uk/documents/6505/FCBU037\\_vol1.pdf](https://www.forestresearch.gov.uk/documents/6505/FCBU037_vol1.pdf)
- Benzian, B. (1967) Manuring young conifers: experiments in some English nurseries. In: *Proceedings Cloquum on Forest Fertilization*, Jyväskylä, Finland:142-170.  
<https://www.forestresearch.gov.uk/documents/6514/FCBU043.pdf>
- Benzian B, Smith HA (1973) Nutrient concentrations of healthy seedlings and transplants of *Picea sitchensis* and other conifers grown in English forest nurseries. *Forestry* 46(1): 55-69.  
<https://doi.org/10.1093/forestry/46.1.55>
- Berry CR (1980) Sewage sludge effects soil properties and growth of slash pine seedlings in a Florida nursery. In: *Proceedings: Lantz CW (ed) Southern Nursery Conference*. USDA Forest Service, State and Private Forestry, Atlanta, GA: 46-51.

- [https://rngr.net/publications/proceedings/1980/sewage-sludge-affects-soil-properties-and-growth-of-slash-pine-seedlings-in-a-florida-nursery/at\\_download/file](https://rngr.net/publications/proceedings/1980/sewage-sludge-affects-soil-properties-and-growth-of-slash-pine-seedlings-in-a-florida-nursery/at_download/file)
- Berenyi NM, Brenneman BB, Owens EG (1971) The effect of magnesium on the growth of loblolly pine grown in two sandhills soils in the greenhouse. Research Report 61. Westvaco, Forest Research. 6 p. <https://d.lib.ncsu.edu/collections/catalog/mc00496-002-bx0011-002-061>
- Berenyi NM, Brenneman BB, Owens EG (1972) 1971 nutrient elements greenhouse study with loblolly pine. Research Report 65. Westvaco, Forest Research. 9 p. <https://d.lib.ncsu.edu/collections/catalog/mc00496-002-bx0011-002-065>
- Binns WO, Mayhead GJ, MacKenzie JM (1980) Nutrient deficiencies of conifers in British forests. An illustrated guide. Leaflet 76. Forestry Commission. 23 p. [https://www.forestresearch.gov.uk/documents/7845/FCLF076\\_1980.pdf](https://www.forestresearch.gov.uk/documents/7845/FCLF076_1980.pdf)
- Blackmon BG (1969) Response of loblolly pine (*Pinus taeda* L.) seedlings to various levels and combinations of nitrogen and phosphorus. Baton Rouge, LA: Louisiana State University. 164 p. Ph.D. dissertation. [https://digitalcommons.lsu.edu/cgi/viewcontent.cgi?article=2637&context=gradschool\\_disstheses](https://digitalcommons.lsu.edu/cgi/viewcontent.cgi?article=2637&context=gradschool_disstheses)
- Bolton J, Benzian B (1970) Sulphur as a nutrient for Sitka spruce (*Picea sitchensis*) seedlings and radish (*Raphanus sativus*) grown on a sandy podzol in England. The Journal of Agricultural Science 74(3): 501-504. <https://doi.org/10.1017/S0021859600017615>
- Boxman AW, Krabbendam H, Bellemakers MJ, Roelofs JG (1991) Effects of ammonium and aluminium on the development and nutrition of *Pinus nigra* in hydroculture. Environ Pollut 73(2): 119-136. [https://doi.org/10.1016/0269-7491\(91\)90018-R](https://doi.org/10.1016/0269-7491(91)90018-R)
- Boyer JN, South DB (1985) Nutrient content of nursery-grown loblolly pine seedlings. Circular 282. Alabama Agricultural Experiment Station, Auburn University, Auburn University, AL: 27 p. <http://131.204.73.195/bitstream/handle/11200/2067/1279CIRC.pdf>
- Boynton D, Burrell AB (1944) Potassium-induced magnesium deficiency in the McIntosh apple tree. Soil Sci 58(6): 441-454. [https://journals.lww.com/soilsci/Citation/1944/12000/Potassium\\_Induced\\_Magnesium\\_Deficiency\\_in\\_the.4.aspx](https://journals.lww.com/soilsci/Citation/1944/12000/Potassium_Induced_Magnesium_Deficiency_in_the.4.aspx)
- Bridger GL, Salutsky ML, Starostka RW (1962) Micronutrient sources, metal ammonium phosphates as fertilizers. J Agr Food Chem 10(3): 181-188. <https://doi.org/10.1021/jf60121a006>
- Briggs RD (2008) Soils and nutrition: A forest nursery perspective. In: Dumroese RK; Riley LE, tech. coords. National Proceedings: Forest and Conservation Nursery Associations-2007. RMRS-P-57. USDA, Forest Service, Rocky Mountain Research Station, Fort Collins, CO: 55-64. [https://rngr.net/publications/proceedings/2007/soils-and-nutrition-a-forest-nursery-perspective/at\\_download/file](https://rngr.net/publications/proceedings/2007/soils-and-nutrition-a-forest-nursery-perspective/at_download/file)
- Brix H, van den Driessche R (1974) Mineral nutrition of container grown tree seedlings. In: Tinus RW, Stein WI, Balmer WE (eds) Proceedings, North American Containerized Forest Tree Seedling Symposium. Publication 68. Great Plains Agric. Council. Denver, CO: 77-84. <https://babel.hathitrust.org/cgi/pt?id=mdp.39015006879780&view=1up&seq=7>
- Bryson GM, Mills HA (eds) (2014) Plant analysis handbook IV. Micro-Macro Publishing: Athens, Georgia. 600 p. [https://www.researchgate.net/publication/271849765\\_Plant\\_Analysis\\_Handbook\\_IV](https://www.researchgate.net/publication/271849765_Plant_Analysis_Handbook_IV)
- Bryson HL (1980) *Pisolithus tinctorius* mycobiont inoculations as a factor in performance of containerized and bare-root shortleaf pine seedlings on lignite minesoils in Panola County, Texas. DF thesis, Stephen F Austin State University, Nacogdoches. 418 p. <https://scholarworks.sfasu.edu/cgi/viewcontent.cgi?article=1011&context=etds>
- Bueno SW (1991) Analysis of soil chemical properties at Saratoga Forest Tree Nursery. PhD thesis, State University of New York, Syracuse. 118 p.
- Bunting WR (1980) Seedling quality: growth and development—soil relationships, seedling growth and development, density control relationships. In: Proceedings of the North American Forest Tree Nursery Soils Workshop. Syracuse, NY: 21-42. <https://babel.hathitrust.org/cgi/pt?id=pst.000007190872&view=1up&seq=9&skin=2021>

- Cakmak I, Yazici AM (2010) Magnesium: a forgotten element in crop production. *Better crops* 94(2): 23-25. [article-201006-better-crops-magnesium.pdf](https://doi.org/10.1006-better-crops-magnesium.pdf) ([ks-minerals-and-agriculture.com](http://ks-minerals-and-agriculture.com))
- Carlson LW (1979) Guidelines for rearing containerized conifer seedlings in the Prairie Provinces. Information Report NOR-X-214. Environment Canada. Northern Forest Research Centre. Edmonton, Alberta: 62 p. <https://cfs.nrcan.gc.ca/publications?id=11650>
- Carter DR, Allen HL, Fox TR, Albaugh TJ, Rubilar RA, Campoe OC, Cook RL (2021) A 50-year retrospective of the Forest Productivity Cooperative in the southeastern United States: regionwide trials. *J Forest* 119(1): 73-85. <https://doi.org/10.1093/jofore/fvaa046>
- Chaganti VN, Culman SW, Herms C, Sprunger CD, Brock C, Soto AL, Doohan D (2021) Base cation saturation ratios, soil health, and yield in organic field crops. *Agron J* 113(5): 4190-4200. <https://doi.org/10.1002/agj2.20785>
- Chen HF (1956) Effects of peat additions and seedling density upon development and chemical composition of Douglas-fir (*Pseudotsuga Menziesii*) nursery stock. MF thesis, University of Washington, Seattle. 34 p. <http://thuir.thu.edu.tw/retrieve/16905/V.2+n.2%2821-34%29.pdf>
- Christersson L (1973) The effect of inorganic nutrients on water economy and hardness of conifers. 1. The effect of varying potassium, calcium, and magnesium levels on water content, transpiration rate, and the initial phase of development of frost hardness of *Pinus sylvestris* L. seedlings. *Studia Forestalia Suecica* 103: 1-28. <https://pub.epsilon.slu.se/5717/1/SFS103.pdf>
- Classen HG, Schimatschek HF, Wink K (2004) Magnesium in human therapy. *Metal ions in biological systems* 41: 41-70. [https://www.academia.edu/download/51660916/Astrid\\_Sigel\\_Helmut\\_Sigel\\_Metal\\_Ions\\_and\\_Their\\_Complexes\\_in\\_Medication.pdf#page=93](https://www.academia.edu/download/51660916/Astrid_Sigel_Helmut_Sigel_Metal_Ions_and_Their_Complexes_in_Medication.pdf#page=93)
- Coleman M, Dunlap I, Dutton D, Bledsoe C (1987) Nursery and field evaluation of compost-grown conifer seedlings. *Tree Planters' Notes* 38(2): 22-27. [https://rngr.net/publications/tpn/38-2/nursery-and-field-evaluation-of-compost-grown-conifer-seedlings/at\\_download/file](https://rngr.net/publications/tpn/38-2/nursery-and-field-evaluation-of-compost-grown-conifer-seedlings/at_download/file)
- Cotton DR (1964) The influence of soil characteristics and fertilizer treatment on growth and chemical composition of *Pinus resinosa*. MS thesis. McGill University, Montreal. 109 P. <https://escholarship.mcgill.ca/downloads/qv33s111v>
- Crowther EM (1950) Nutritional problems in forest nurseries. Report for 1949, Rothamsted Research: 122-129. <https://doi.org/10.23637/ERADOC-1-71>
- Culman S, Mann M, Sharma S, Saeed M, Fulford A, Lindsey L, Joern B (2019) Converting between Mehlich-3, Bray P, and ammonium acetate soil test values. ANR-75, The Ohio State University, Ohio. 4 p. <https://www.canr.msu.edu/soilfertility/Files/Bulletins/Bray%20to%20Mehlich%20conversion.pdf>
- Cumming JR, Weinstein LH (1990) Aluminum-mycorrhizal interactions in the physiology of pitch pine seedlings. *Plant Soil* 125(1): 7-18. <https://doi.org/10.1007/BF00010739>
- Danielson RM (1966) The effect of soil fumigation on seedling growth, mycorrhizae and the associated microflora of loblolly pine (*Pinus taeda* L.) roots. MS thesis, North Carolina State University, Raleigh. 148 p.
- Davey CB (2002) Using soil test results to determine fertilizer applications. In: Dumroese RK, Riley LE, Landis TD (eds) *Proceedings, forest and conservation nursery associations-1999, 2000, and 2001*. RMRS-P-24. USDA Forest Service, Rocky Mountain Research Station, Ogden UT: 22-26. <https://rngr.net/publications/proceedings/1999/davey.pdf>
- Davey CB, McNabb K (2019) The management of seedling nutrition. In: McNabb K, Pike C (eds) *Nursery Guide for the Production of Bareroot Hardwood Seedlings*. Agriculture Handbook 733. USDA Forest Service, Washington, DC: 75-87. [https://rngr.net/publications/a-nursery-guide-for-the-production-of-bareroot-hardwood-seedlings/introduction/at\\_download/file](https://rngr.net/publications/a-nursery-guide-for-the-production-of-bareroot-hardwood-seedlings/introduction/at_download/file)
- Davis AS, Jacobs DF, Wightman KE (2007a) Organic matter amendment of fallow forest tree seedling nursery soils influences soil properties and biomass of a sorghum cover crop. *Tree Planter's Notes* 52(1): 4-8. [https://rngr.net/publications/tpn/52-1/organic-matter-amendment-of-fallow-forest-tree-seedling-nursery-soils-influences-soil-properties-and-biomass-of-a-sorghum-cover-crop/at\\_download/file](https://rngr.net/publications/tpn/52-1/organic-matter-amendment-of-fallow-forest-tree-seedling-nursery-soils-influences-soil-properties-and-biomass-of-a-sorghum-cover-crop/at_download/file)

- Davis M, Zue J, Clinton P (2015) Planted-forest nutrition. NZ For Res Inst Repot Info, Sheet 126 p. [https://www.researchgate.net/profile/Jianming-Xue/publication/286931750\\_Planted\\_forest\\_Nutrition/links/5670f60508ae2b1f87acfb65/Planted-forest-Nutrition.pdf](https://www.researchgate.net/profile/Jianming-Xue/publication/286931750_Planted_forest_Nutrition/links/5670f60508ae2b1f87acfb65/Planted-forest-Nutrition.pdf)
- Davis MR, Coker G, Parfitt RL, Simcock R, Clinton PW, Garrett LG, Watt MS (2007b) Relationships between soil and foliar nutrients in young densely planted mini-plots of *Pinus radiata* and *Cupressus lusitanica*. *Forest EcolManag* 240(1-3): 122-130. [https://www.academia.edu/download/45502190/Relationships\\_between\\_soil\\_and\\_foliar\\_nu20160510-32745-1h6icqg.pdf](https://www.academia.edu/download/45502190/Relationships_between_soil_and_foliar_nu20160510-32745-1h6icqg.pdf)
- Deines J (1973) The effects of fertilization on the growth and development of 1-0 sycamore (*Platanus occidentalis* L.), sweetgum (*Liquidambar styraciflua* L.) and green ash (*Fraxinus pennsylvanica* Marsh.) seedlings. PhD thesis, North Carolina State University, Raleigh. 80 p
- Deshmukh J, Ray S (2019) Effectiveness of application of hot water with Epsom salt v/s plain hot water on knee joint pain among geriatric women. *The Pharma Innovation Journal* 8(6): 434-441. <https://www.thepharmajournal.com/archives/2019/vol8issue6/PartH/8-5-169-652.pdf>
- Devitt DA, Morris RL, Fenstermaker LK (2005) Foliar damage, spectral reflectance, and tissue ion concentrations of trees sprinkle irrigated with waters of similar salinity but different chemical composition. *HortSci* 40(3): 819-826. <https://journals.ashs.org/hortsci/downloadpdf/journals/hortsci/40/3/article-p819.pdf>
- Dickson A, Leaf AL, Hosner JF (1960) Seedling quality - soil fertility relationships of white spruce, and red and white pine in nurseries. *The Forestry Chronicle* 36(3): 237-241. <https://pubs.cif-ific.org/doi/pdf/10.5558/tfc36237-3>
- Dobrahner J, Lowery B, Iyer JG (2004) Slow-release fertilization reduces nitrate leaching in bareroot production of *Pinus strobus* seedlings. In: Riley LE, Dumroese RK, Landis TD. (eds). National proceedings: Forest and Conservation Nursery Associations. RMRS-P-33. USDA Forest Service, Rocky Mountain Research Station. Fort Collins, CO: 129-139. [https://rngr.net/publications/proceedings/2003/PDF.2004-06-08.4503/at\\_download/file](https://rngr.net/publications/proceedings/2003/PDF.2004-06-08.4503/at_download/file)
- Donald DGM (1991) Nursery fertilization of conifer planting stock. In: van den Driessche R (ed) *Mineral Nutrition of Conifer Seedlings*: 135-167.
- Donald DGM, Young I (1982) The growth of pine seedlings in South African forest nurseries. *South African Forestry Journal* 123(1): 36-50. <https://doi.org/10.1080/00382167.1982.9628852>
- Dumbroff EB, Michel BE (1967) The expression of interionic relationships in *Pinus elliotii*. *Plant Physiol* 42(11): 1465-1471. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1086752/pdf/plntphys00503-0003.pdf>
- Edwards NT, Taylor GE, Adams MB, Simmons GL, Kelly JM (1990) Ozone, acidic rain and soil magnesium effects on growth and foliar pigments of *Pinus taeda* L. *Tree Physiol* 6(1): 95-104. <https://doi.org/10.1093/treephys/6.1.95>
- Engstrom HE, Stoeckeler JH (1941) Nursery practice for trees and shrubs suitable for planting on the prairie-plains. Miscellaneous publication 434. USDA. Washington, DC: 159 p. <https://babel.hathitrust.org/cgi/pt?id=mdp.39015000753452&view=1up&seq=3&skin=2021>
- Flaten CM (1939) A study on chlorosis of *Pinus resinosa* in a forest nursery, with special reference to some mineral deficiencies. MF thesis. University of Michigan, Ann Arbor: 35 p. <https://deepblue.lib.umich.edu/bitstream/handle/2027.42/114430/39015003261644.pdf?sequence=2>
- Flinn DW, Homans P, Craig FG (1980) Survey of the nutrient status of *Pinus radiata* seedlings and of soil properties in three Victorian nurseries. *Aust Forestry* 43(1): 58-66. <https://doi.org/10.1080/00049158.1980.10674246>
- Fox WF (1904) *Forest nurseries and nursery methods in Europe*. JB Lyon Company, Albany. <https://babel.hathitrust.org/cgi/pt?id=loc.ark:/13960/t68348s41&view=1up&seq=3&skin=2021>
- Franklin JA, Zwiazek JJ, Renault S, Croser C (2002) Growth and elemental composition of jack pine (*Pinus banksiana*) seedlings treated with sodium chloride and sodium sulfate. *Trees* 16(4): 325-330.

- <https://www.academia.edu/download/54065916/s00468-002-0175-520170805-15730-18v93l9.pdf>
- Gaspar AP, Laboski CAM (2016) Base saturation: What is it? Should I be concerned? Does it affect my fertility program? In: Proceedings Wis. Crop Manage. Conf 5: 55-61.  
<https://extension.soils.wisc.edu/wp-content/uploads/sites/68/2014/02/WCMC-2016-Complete-Proceedings.pdf#page=63>
- Giertych MM, Farrar JL (1961) The effect of photoperiod and nitrogen on the growth and development of seedlings of jack pine. *Can J Bot* 39(5): 1247-1254. <https://doi.org/10.1139/b61-109>
- Gleason JF (1988) Fertilization of 2-0 ponderosa pine seedlings in the nursery and field: morphology, physiology, and field performance. MS thesis, Oregon State University, Corvallis. 109 p.  
<https://ir.library.oregonstate.edu/downloads/5712mb473>
- Goodrich BA, Jacobi WR (2012) Foliar damage, ion content, and mortality rate of five common roadside tree species treated with soil applications of magnesium chloride. *Water, Air, & Soil Pollution* 223(2): 847-862. <https://doi.org/10.1007/s11270-011-0907-5>
- Goslin WE (1959) Effects of deficiencies of essential elements on the development and mineral composition of seedlings of Scots pine (*Pinus sylvestris* L.). PhD thesis, The Ohio State University. Columbus. 114 p. [https://etd.ohiolink.edu/pg\\_10?::NO:10:P10\\_ETD\\_SUBID:123562](https://etd.ohiolink.edu/pg_10?::NO:10:P10_ETD_SUBID:123562)
- Grzebisz W (2015) Magnesium. In: Barker AV, Pilbeam DJ (eds) *Handbook of plant nutrition*. Taylor & Francis, Boca Raton, FL: 199-260. <https://doi.org/10.1201/9781420014877>
- Hacskaylo J, Finn RF, Vimmerstedt JP (1969) Deficiency symptoms of some forest trees. *Research Bulletin* 1015. Ohio Agricultural Research and Development Center, Wooster, OH: 69 p.  
[https://kb.osu.edu/bitstream/handle/1811/62823/1/OARDC\\_research\\_bulletin\\_n1015.pdf](https://kb.osu.edu/bitstream/handle/1811/62823/1/OARDC_research_bulletin_n1015.pdf)
- Hallett RD (1980) Experience in the use of soil analysis data. In: Proceedings of the North American Forest Tree Nursery Soils Workshop. Syracuse, NY: 296-298.  
<https://babel.hathitrust.org/cgi/pt?id=pst.000007190872&view=1up&seq=9&skin=2021>
- Hans RR (2013) Initial growth responses to controlled release fertilizer application at establishment of commercial forestry species in South Africa. PhD thesis, Stellenbosch University. Stellenbosch. 158 p. <http://scholar.sun.ac.za/handle/10019.1/85609>
- Hardy DH, Tucker MR, Stokes C (2013) Understanding the soil test report. Miscellaneous Publication. NC Department of Agriculture. 9 p. <https://www.ncagr.gov/agronomi/pdf/ustr.pdf>
- Hart PBS, Widdowson JP, Watts HM, Chu-Chou M (1980) Response of *Pinus caribaea* var. *hondurensis* seedlings to mycorrhizal inoculum, phosphorus and pH. *Australian Forest Research* 10(4): 389-396. [https://drive.google.com/file/d/0B9Klbbp1-PmyVF80M0YwcEtta1U/view?resourcekey=0-YAUqNKUikuxDPw\\_Rqv8TEg](https://drive.google.com/file/d/0B9Klbbp1-PmyVF80M0YwcEtta1U/view?resourcekey=0-YAUqNKUikuxDPw_Rqv8TEg)
- Hart PBS, Widdowson JP (1981) The response of caribbean pine, green panic, and siratro to fertiliser on soils of the 'Eua Uplands, Tonga. *New Zealand Journal of Experimental Agriculture* 9(3-4): 255-262. <https://doi.org/10.1080/03015521.1981.10425423>
- Hauer-Jákli M, Tränkner M (2019) Critical leaf magnesium thresholds and the impact of magnesium on plant growth and photo-oxidative defense: a systematic review and meta-analysis from 70 years of research. *Front Plant Sci* 10: 766.  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6592071/>
- Heckman JR, Sims JT, Beegle DB, Coale FJ, Herbert SJ, Bruulsema TW, Bamka WJ (2003) Nutrient removal by corn grain harvest. *Agron J* 95(3): 587-591.  
<https://doi.org/10.2134/agronj2003.5870>
- Hinesley LE, Maki TE (1980) Fall fertilization helps longleaf pine nursery stock. *Southern Journal of Applied Forestry* 4(3): 132-135. <https://doi.org/10.1093/sjaf/4.3.132>
- Hobbs CH (1944) Studies on mineral deficiency in pine. *Plant Physiol* 19(4): 590-602.  
<https://doi.org/10.1104/pp.19.4.590>
- Hopmans P, Flinn DW (1983) Nutrient requirements in three Victorian radiata pine nurseries with contrasting soils. *Aust Forestry* 46(2): 111-117.  
<https://doi.org/10.1080/00049158.1983.10674386>
- Howell J (1932) The Effect of the concentration of the culture solution on seedlings of ponderosa pine. *J Forest* 30(7): 829-830. <https://doi.org/10.1093/jof/30.7.829>

- Huberman MA (1935) Illustrated summary of Stuart Forest Nursery practice and research. Forest History Society, Durham, NC:175 p. <https://foresthistor.org/wp-content/uploads/2017/01/Illustrated-History-of-Stuart-Nursery.pdf>
- Hunter IR (1996) The occurrence and treatment of magnesium deficiency in radiata pine in New Zealand. New Zealand Forest Research Institute, FRI Bulletin No. 172. 136 p. <https://scion.contentdm.oclc.org/digital/api/collection/p20044coll6/id/166/download>
- Hunter IR, Prince JM, Graham JD, Nicholson GM (1986) Growth and nutrition of *Pinus radiata* on rhyolitic tephra as affected by magnesium fertiliser. NZ J Forestry Sci 16(2): 152-165. <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.714.6809>
- Ingestad T (1960) Magnesium deficiency in nurseries of forest trees. Vaxt-Narings-Nytt 16(3): 30-32.
- Irwin KM, Duryea ML, Stone EL (1998) Fall-applied nitrogen improves performance of 1-0 slash pine nursery seedlings after outplanting. South J Appl For 22(2): 111-116. <https://doi.org/10.1093/sjaf/22.2.111>
- Iyer JG (1965) Effect of Dacthal 75 on the growth of nursery stock. Tree Planters' Notes 71: 13-16. [https://rngr.net/publications/tpn/16-2/pdf.2005-05-09.4763443573/at\\_download/file](https://rngr.net/publications/tpn/16-2/pdf.2005-05-09.4763443573/at_download/file)
- Iyer JG, Schulte EE, Randall GW (1971) Relationship between foliar composition of red-pine and jack-pine seedlings and vulnerability to *Lophodermium* needle-cast disease. Plant Soil 35(1): 213-215. <https://link.springer.com/content/pdf/10.1007/BF01372651.pdf>
- Iyer JG, Dobrahner J, Lowery B, VandeHey J (2002) slow-release fertilizers in bareroot nurseries. In: Dumroese RK, Riley LE, Landis TD (eds) Proceedings, forest and conservation nursery associations-1999, 2000, and 2001. RMRS-P-24. USDA Forest Service, Rocky Mountain Research Station, Ogden UT: 112-110. <https://rngr.net/publications/proceedings/2000/iyer%2Cdobrahner.pdf>
- Januszek K, Stępniewska H, Błońska E, Molicka J, Kozieł K, Gdula A, Wójs A (2014) Impact of aluminum sulphate fertilizer on selected soil properties and the efficiency and quality of pine seedlings in the forest ground tree nursery. LeśnePraceBadawcze 75(2): 127-138. <https://depot.ceon.pl/bitstream/handle/123456789/5193/doi-10-2478-frp-2014-0012%20e.pdf?sequence=1&isAllowed=y>
- Jokela EJ (2004) Nutrient management of southern pines. In: Dickens ED, Barnett JP, Hubbard WG, Jokela EJ (eds) Slash pine: still growing and growing. GTR-SRS-76. USDA Forest Service, Southern Research Station, Asheville, NC: 27-35. [https://www.srs.fs.usda.gov/pubs/gtr/gtr\\_srs076.pdf](https://www.srs.fs.usda.gov/pubs/gtr/gtr_srs076.pdf)
- Kelly JM, Barber SA (1991) Magnesium uptake kinetics in loblolly pine seedlings. Plant Soil 134(2): 227-232. <https://link.springer.com/article/10.1007/BF00012040>
- Knight PJ (1978a) Fertilizer practice in New Zealand forest nurseries. NZJ For Sci 8(1): 27-53. [https://www.scionresearch.com/\\_data/assets/pdf\\_file/0018/59022/NZJFS811978KNIGHT27\\_53.pdf](https://www.scionresearch.com/_data/assets/pdf_file/0018/59022/NZJFS811978KNIGHT27_53.pdf)
- Knight PJ (1978b) The nutrient content of *Pinus radiata* seedlings: a survey of planting stock from 17 New Zealand forest nurseries. NZJ For Sci 8(1): 54-69. [https://www.scionresearch.com/\\_data/assets/pdf\\_file/0006/37185/NZJFS811978KNIGHT54\\_69.pdf](https://www.scionresearch.com/_data/assets/pdf_file/0006/37185/NZJFS811978KNIGHT54_69.pdf)
- Knight PJ (1981) The maintenance of productivity in forest nurseries. In: FRI Symposium 22. New Zealand Forest Service, Forest Research Institute: 48-69.
- Koll PJ (2009) Effects of conifer sawdust, hardwood sawdust, and peat on soil properties and bareroot conifer seedlings development. MS thesis, Michigan Technological University, Houghton. 42 p. <https://digitalcommons.mtu.edu/cgi/viewcontent.cgi?article=1139&context=etds>
- Kopittke PM, Menzies NW (2007) A review of the use of the basic cation saturation ratio and the "ideal" soil. Soil Sci Soc Am J 71(2): 259-265. <http://www.agvise.com/wp-content/uploads/2015/02/SSSA-Cation-Ratios.pdf>
- Kormanik PP, Sung SJS, Kormanik TL (1994) Irrigating and fertilizing to grow better nursery seedlings. In: Landis TD (ed). Proceedings: northeastern and intermountain forest and conservation nursery associations. GTR-RM-243. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO: 115-121. [https://rngr.net/publications/proceedings/1993/kormanik.pdf/at\\_download/file](https://rngr.net/publications/proceedings/1993/kormanik.pdf/at_download/file)

- Landis TD (1988) Management of forest nursery soils dominated by calcium salts. *New Forest* 2(3): 173-193. <https://doi.org/10.1007/BF00029987>
- Landis TD (1996) Secondary nutrients - magnesium. *Forest Nursery Notes* 16(2): 5-8. [https://rngr.net/publications/fnn/1996-summer/articles/secondary-nutrients-magnesium/at\\_download/file](https://rngr.net/publications/fnn/1996-summer/articles/secondary-nutrients-magnesium/at_download/file)
- Landis TD, Tinus RW, McDonald SE, Barnett JP (1989) Seedling nutrition and irrigation. In: *The container tree nursery manual*. Agricultural Handbook 674, Volume 4. USDA Forest Service, Washington, DC: 119 p. <https://rngr.net/publications/ctnm/volume-4>
- Landis TD, Pinto JR, Davis AS (2009) Fertigation-injecting soluble fertilizers into the irrigation system. *Forest Nursery Notes* 29(2): 4-13. [https://www.fs.fed.us/rm/pubs\\_other/rmrs\\_2009\\_landis\\_t002.pdf](https://www.fs.fed.us/rm/pubs_other/rmrs_2009_landis_t002.pdf)
- Larsen HS, South DB, Boyer JN (1988) Foliar nitrogen content at lifting correlates with early growth of loblolly pine seedlings from 20 nurseries. *South J Appl For* 12(3): 181-185. <https://doi.org/10.1093/sjaf/12.3.181>
- Leach GN, Gresham HH (1983) Early field performance of loblolly pine seedlings with *Pisolithus tinctorius* ectomycorrhizae on two lower coastal plain sites. *South J Appl For* 7(3): 149-153. <https://academic.oup.com/sjaf/article-abstract/7/3/149/4794396>
- Leaf, AL (1968) K, Mg, and S deficiencies in forest trees. In: *Forest Fertilization-Theory and Practice*. Tennessee Valley Authority, Muscle Shoals, AL: 88-122. <https://catalog.hathitrust.org/Record/001516717>
- Leski T, Aučina A, Skridaila A, Pietras M, Riepšas E, Rudawska M (2010) Ectomycorrhizal community structure of different genotypes of Scots pine under forest nursery conditions. *Mycorrhiza* 20(7): 473-481. <https://doi.org/10.1007/s00572-010-0298-2>
- Lipman CB (1916) A critique of the hypothesis of the lime-magnesia ratio. *The Plant World* 19(4): 83-105. <https://www.jstor.org/stable/pdf/43477507.pdf>
- Lunt HA (1938) The use of fertilizers in the coniferous nursery. Bulletin 416, Connecticut Agricultural Experiment Station: New Haven CT: 721-766. <https://portal.ct.gov/-/media/CAES/DOCUMENTS/Publications/Bulletins/B416pdf.pdf>
- Lyle ES (1969) Mineral deficiency symptoms in loblolly pine seedlings. *Agron J* 61(3): 395-398. <https://doi.org/10.2134/agronj1969.00021962006100030019x>
- Lyle ES, Adams F (1971) Effect of available soil calcium on taproot elongation of loblolly pine (*Pinus taeda* L.) seedlings. *Soil Sci Soc Am J* 35(5): 800-805. <https://doi.org/10.2136/sssaj1971.03615995003500050046x>
- Lyle ES, Pearce ND (1968) Sulfur deficiency in nursery seedlings may be caused by concentrated fertilizers. *Tree Planters' Notes* 19(1): 9-10. [https://rngr.net/publications/tpn/19-1/pdf.2005-05-20.5572607495/at\\_download/file](https://rngr.net/publications/tpn/19-1/pdf.2005-05-20.5572607495/at_download/file)
- Madgwick HAI, Ovington JD (1959) The chemical composition of precipitation in adjacent forest and open plots. *Forestry* 32(1): 14-22. <https://doi.org/10.1093/forestry/32.1.14>
- Madgwick HAI (1975) Branch growth of *Pinus resinosa* Ait. with particular reference to potassium nutrition. *Can J Forest Res* 5(4): 509-514. <https://doi.org/10.1139/x75-074>
- Majid NM (1984) Some aspects of boron, copper and iron nutrition of lodgepole pine and Douglas-fir. PhD thesis, University of British Columbia. Vancouver. 172 p. <https://open.library.ubc.ca/collections/831/831/items/1.0096652>
- Maki TE, Henry BW (1951) Root-rot control and soil improvement at the Ashe Forest Nursery. Occasional Paper 119. USDA Forest Service, Southern Forest Experiment Station. New Orleans, LA: 23 p. <https://archive.org/details/CAT31363714/page/16/mode/2up>
- Malavolta E, Sarruge JR, Haag HP, Vencovsky R, Santos CFO, Valsechi O, Scoton LC, Coelho RSG (1970) The relation of the concentration of macronutrients in the substrate and in the foliage to cell wall thickness and cellulose concentration in the xylem of slash pine (*Pinus elliottii*). *Anais da Escola Superior de Agricultura Luiz de Queiroz* 27: 295-333. <https://www.scielo.br/pdf/aesalq/v27/22.pdf>
- Manikam D, Srivastava PBL (1980) The growth response of *Pinus caribaea* mor. var. *hondurensis* bar and golf seedlings to fertilizer application on the Serdang soil series. *Forest EcolManag* 3: 127-139.



- [https://doi.org/10.1016/0378-1127\(80\)90010-9](https://doi.org/10.1016/0378-1127(80)90010-9)
- Martian, BF (1989) Soil management practices at the Big Sioux Nursery. In: Landis TD (ed) Proceedings, Intermountain Forest Nursery Association. GTR RM-184. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station: 82-85.  
[https://rngr.net/publications/proceedings/1989/martian.pdf/at\\_download/file](https://rngr.net/publications/proceedings/1989/martian.pdf/at_download/file)
- Marx DH (1990) Soil pH and nitrogen influence *Pisolithus* ectomycorrhizal development and growth of loblolly pine seedlings. *Forest Sci* 36 (2): 224-245.  
<https://doi.org/10.1093/forestscience/36.2.224>
- Marx DH, Cordell CE, Kenney DS, Mexal JG, Artman JD, Riffle JW, Molina RJ (1984) Commercial vegetative inoculum of *Pisolithus tinctorius* and inoculation techniques for development of ectomycorrhizae on bare-root tree seedlings. *For Sci* 30(3): Monograph 25.  
[https://academic.oup.com/forestscience/article-abstract/30/suppl\\_1/a0001/4656736](https://academic.oup.com/forestscience/article-abstract/30/suppl_1/a0001/4656736)
- Mason WL, Negussie G, Hollingsworth MK (1995) Seed pretreatments and nursery regimes for raising Macedonian pine (*Pinus peuce* Grisebach). *Forestry* 68(3): 255-264.  
<https://doi.org/10.1093/forestry/68.3.255>
- May JT (1957) Effects of soil management practices in a forest tree nursery on soil properties and on loblolly pine seedlings. PhD thesis, Michigan State University, East Lansing. 215 p.
- May JT, Johnson HH, Gilmore AR (1962) Chemical composition of southern pine seedlings. Research Paper 10. Georgia Forest Research Council, Macon, GA: 11 p.
- May JT (1984) Nutrients and fertilization. In: Lantz CW ed: Southern Pine Nursery Handbook. Atlanta, GA: USDA Forest Service, Southern Region: 1201-1241.  
[https://rngr.net/publications/spnh/PDF.2003-08-12.3705/at\\_download/file](https://rngr.net/publications/spnh/PDF.2003-08-12.3705/at_download/file)
- Maxwell JW (1988) Macro and micronutrient programmes in B.C. bareroot nurseries. In: Landis TD (ed) Proceedings, Combined Meeting of the Western Forest Nursery Associations. GTR-RM-167. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO: 11-14. [https://rngr.net/publications/proceedings/1988/maxwell.pdf/at\\_download/file](https://rngr.net/publications/proceedings/1988/maxwell.pdf/at_download/file)
- McConnell RC, Klages MG (1969) Forest nursery soils of northern Idaho and western Montana. Montana Agricultural Experiment Station Montana State University, Bozeman. 33 p. <https://babel.hathitrust.org/cgi/pt?id=uiug.30112019700563&view=1up&seq=6&skin=2021>
- McDaniel VE (1931) Nursery practice on the Oregon forest nursery. MS thesis. Oregon State University, Corvallis. 52 p. <http://hdl.handle.net/1957/16213>
- McNabb K, Heidbreder-Olson E (1998) Results of the 1996 irrigation water quality survey. Research report 98-05. Auburn, AL: Auburn University, Southern Forest Nursery Management Cooperative. 7 p.
- Mellert KH, Göttelein A (2012) Comparison of new foliar nutrient thresholds derived from van den Burg's literature compilation with established central European references. *Eur J Forest Res* 131(5): 1461-1472. <https://link.springer.com/content/pdf/10.1007/s10342-012-0615-8.pdf>
- Menzies MI, Holden DG, Klomp BK (2001) Recent trends in nursery practice in New Zealand. *New Forest* 22(1): 3-17. <https://doi.org/10.1023/A:1012027013173>
- Metson AJ (1974) I. Some factors governing the availability of soil magnesium: A review, *New Zealand Journal of Experimental Agriculture* 2:3: 277-319.  
<https://doi.org/10.1080/03015521.1974.10427689>
- Mexal JG, Fisher JT (1987) Organic matter amendments to a calcareous forest nursery soil. *New Forest* 1(4): 311-323. <https://doi.org/10.1007/BF00031741>
- Miller SP, Cumming JR (2000) Effects of serpentine soil factors on Virginia pine (*Pinus virginiana*) seedlings. *Tree Physiol* 20(16): 1129-1135. <https://doi.org/10.1093/treephys/20.16.1129>
- Mitchell AD (2000) Magnesium fertiliser effects on forest soils under *Pinus radiata*. PhD thesis. Massey University, Palmerston North. 247 p.  
[https://mro.massey.ac.nz/bitstream/handle/10179/2129/02\\_whole.pdf?sequence=1&isAllowed=y](https://mro.massey.ac.nz/bitstream/handle/10179/2129/02_whole.pdf?sequence=1&isAllowed=y)
- Moffat AJ (1994) Nursery sterilization and inoculation regimes for alder production. *Forestry* 67(4): 313-327. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.904.245&rep=rep1&type=pdf>
- Möller A (1904) Karenzerscheinungen bei der Kiefer. *Zeitschrift für Forst- und Jagdwesen* 36: 745-756.

- <https://play.google.com/books/reader?id=GQEWAQAIAAJ&pg=GBS.PA756&hl=en>  
Morrison IK (1974) Mineral nutrition of conifers with special reference to nutrient status interpretation: a review of literature. Publication 1343. Environment Canada, Forestry Canada. Great Lakes Forest Research Centre, Sault Ste. Marie, Ontario: 79 p.  
<https://cfs.nrcan.gc.ca/publications?id=37802>
- Morrissey N, O'Reilly C (2002) Effect of root wrenching in the nursery on the quality of Japanese larch transplants. *Irish Forestry* 59 (1-2): 2-17.  
<https://journal.societyofirishforesters.ie/index.php/forestry/article/download/9920/9010>
- Moser F (1933) The calcium-magnesium ratio in soils and its relation to plant growth. *Jour Am Soc Agron* 25: 365-377. <https://doi.org/10.2134/agronj1933.00021962002500060001x>
- Munson KR (1982) Decomposition, function, and maintenance of organic matter in a sandy nursery soil. PhD thesis, University of Florida, Gainesville. 96 p.  
<http://file.iflora.cn/fastdfs/group1/M00/64/46/wKhnoF2NwXSAMM8tAD0yVvJ61s1354.pdf>
- Murison WF (1960) Macronutrient deficiency and its effect on coniferous growth. *PhD thesis*. University of British Columbia, Vancouver. 235 p.  
<https://open.library.ubc.ca/collections/831/831/items/1.0106016>
- Mylavarapu RS, Sanchez JF, Nguyen JH, Bartos JM (2002) Evaluation of Mehlich-1 and Mehlich-3 extraction procedures for plant nutrients in acid mineral soils of Florida. *Commun Soil Sci Plan* 33(5-6): 807-820. <https://doi.org/10.1081/CSS-120003067>
- North Carolina State Forest Nutrition Cooperative (NCSFNC) (1991) Descriptive statistics and relationships among soil and foliar characteristics in midrotation loblolly pine plantations. Res. Note 7. College of Forest Resources, North Carolina State University, Raleigh, NC: 29 p.
- Olykan S, Payn T, Beets P, Kimberley M (2001) Magnesium fertilisers affected growth, upper mid-crown yellowing, and foliar nutrients of *Pinus radiata*, and soil magnesium concentration. *New Zealand Journal of Forestry Science* 31(1): 34-50.  
[https://www.scionresearch.com/\\_data/assets/pdf\\_file/0008/59354/03-OLYKAN.pdf](https://www.scionresearch.com/_data/assets/pdf_file/0008/59354/03-OLYKAN.pdf)
- Payn TW, Mead DJ, Will GM, Hunter IR (1995) Magnesium nutrition and dry matter allocation patterns in *Pinus radiata*. *New Zealand Journal of Forestry Science* 25(1): 39-48.  
[https://www.scionresearch.com/\\_data/assets/pdf\\_file/0020/17372/NZJFS2511995PAYN39-48.pdf](https://www.scionresearch.com/_data/assets/pdf_file/0020/17372/NZJFS2511995PAYN39-48.pdf)
- Pessin LJ (1937) The effect of nutrient deficiency on the growth of longleaf pine seedlings. USDA Forest Service, Southern Forest Experiment Station. New Orleans LA: Occasional Paper 65: 1-7.  
<https://babel.hathitrust.org/cgi/pt?id=mdp.39015003619577&view=1up&seq=325&skin=2021&q1=pessin>
- Purnell HM (1958) Nutritional studies of *Pinus radiata* Don: 1. Symptoms due to deficiencies of some major elements. *Aust Forestry* 22(2): 82-87. <https://doi.org/10.1080/00049158.1958.10675852>
- Rahmani M, Hodges AW, Kiker CF (2004) Compost users' attitudes toward compost application in Florida. *Compost Science & Utilization* 12(1): 55-60.  
<https://doi.org/10.1080/1065657X.2004.10702158>
- Rodríguez-Trejo DA, Duryea ML (2003) Seedling quality indicators in *Pinus palustris* Mill. *Agrociencia* 37(3): 299-307.  
[https://www.researchgate.net/publication/289349591\\_Seedling\\_quality\\_indicators\\_in\\_Pinus\\_palustris\\_Mill](https://www.researchgate.net/publication/289349591_Seedling_quality_indicators_in_Pinus_palustris_Mill)
- Ross NM (1929) Nursery practice under prairie conditions. *The Forestry Chronicle* 5(1): 38-44.  
<https://pubs.cif-ifc.org/doi/pdf/10.5558/tfc5038-1>
- Rowan SJ (1971) Soil fertilization, fumigation, and temperature affect severity of black root rot of slash pine. *Phytopathology* 61: 184-187.  
[https://www.apsnet.org/publications/phytopathology/backissues/Documents/1971Articles/Phyto61n02\\_184.pdf](https://www.apsnet.org/publications/phytopathology/backissues/Documents/1971Articles/Phyto61n02_184.pdf)
- Rowan SJ (1987) Effects of potassium fertilization in the nursery on survival and growth of pine seedlings in the plantation. Georgia Forest Research Paper 68. Georgia Forestry Commission: 9 p. <http://www.gatrees.gov/resources/publications/research-papers/GFRP68.pdf>

- Rowan SJ, Steinbeck K (1977) Seedling age and fertilization affect susceptibility of loblolly pine to fusiform rust. *Phytopathology* 67(24): 242-246.  
[https://www.apsnet.org/publications/phytopathology/backissues/Documents/1977Articles/Phyto67n02\\_242.pdf](https://www.apsnet.org/publications/phytopathology/backissues/Documents/1977Articles/Phyto67n02_242.pdf)
- Rowe DB (1996) Influence of stock plant nitrogen nutrition on mineral nutrient and carbohydrate status, photosynthesis, orthotropic shoot production, and adventitious rooting of stem cuttings from hedged loblolly pine. PhD thesis, North Carolina State University, Raleigh. 180 p.  
<https://www.proquest.com/openview/d38e0286b846d7f64db195f59e252e32/1?pg-origsite=gscholar&cbl=18750&diss=y>
- Sadreika V (1976) Processing soil and plant data print-outs and recommendations. In: Proceedings, Northeastern Area Nurserymen's Conference, Kemptville, Ontario: 27-46.  
[https://rngr.net/publications/proceedings/1976/processing-soil-and-plant-data-print-outs-and-recommendations/at\\_download/file](https://rngr.net/publications/proceedings/1976/processing-soil-and-plant-data-print-outs-and-recommendations/at_download/file)
- Salmon BC, Arnold PW (1963) The uptake of magnesium under exhaustive cropping. *The Journal of Agricultural Science* 61(3): 421-425. <https://core.ac.uk/download/pdf/162100239.pdf>
- Schaedle M (1959) A study of the growth of Douglas fir (*Pseudotsuga menziesii*) seedlings. MS thesis. University of British Columbia, Vancouver. 171 p.  
<https://open.library.ubc.ca/cIRcle/collections/ubctheses/831/items/1.0106130>
- Schenck CA (1907) Biltmore lectures on silviculture. Brandow Printing Company, NY.  
<https://www.biodiversitylibrary.org/item/59977#page/7/mode/1up>
- Šrámek V, Fadrhonsová V, Vortelová L, Lomský B (2012) Development of chemical soil properties in the western Ore Mts. (Czech Republic) 10 years after liming. *Journal of Forest Science* 58(2): 57-66.  
[https://www.agriculturejournals.cz/publicFiles/72\\_2011-JFS.pdf](https://www.agriculturejournals.cz/publicFiles/72_2011-JFS.pdf)
- Schmidt TL (1991) Factors influencing establishment of eastern redcedar (*Juniperus virginiana* L.) on rangeland. PhD thesis. University of Nebraska, Lincoln. 139 p.  
<https://www.proquest.com/docview/303929676?pq-origsite=gscholar&fromopenview=true>
- Schomaker CE (1969) Growth and foliar nutrition of white pine seedlings as influenced by simultaneous changes in moisture and nutrient supply. *Soil Sci Soc Am J* 33(4): 614-618.  
<https://doi.org/10.2136/sssaj1969.03615995003300040035x>
- Schulte EE, Kelling KE (1985) Soil Calcium to Magnesium Ratios--should You be Concerned? A2986. University of Wisconsin, Madison: 4 p.  
<http://corn.agronomy.wisc.edu/Management/pdfs/a2986.pdf>
- Show SB (1930) Forest nursery and planting practice in the California pine region. Circular 92. USDA, Washington, DC: 75 p.  
<https://babel.hathitrust.org/cgi/pt?id=uiug.30112019276846&view=1up&seq=1&skin=2021>
- Slaton SH, Iyer JG (1974) Manganese compounds harmful to planting stock under some soil conditions. *Tree Planters' Notes* 25(2): 19-21. [https://rngr.net/publications/tpn/25-2/PDF.2003-09-25.2347/at\\_download/file](https://rngr.net/publications/tpn/25-2/PDF.2003-09-25.2347/at_download/file)
- Smits MM, Wallander H (2017) Role of mycorrhizal symbiosis in mineral weathering and nutrient mining from soil parent material. In: *Mycorrhizal Mediation of Soil*: 35-46.  
<https://doi.org/10.1016/B978-0-12-804312-7.00003-6>
- Solan, FM, Bickelhaupt DH, Leaf AL (1979) Soil and plant analytical services for tree nurseries. In: Proceedings Northeastern Area Nurseryman's Conference. USDA Forest Service, Northeastern area State and Private Forestry, Broomall, PA: 35-42.  
[https://rngr.net/publications/proceedings/1979/soil-and-plant-analytical-services-for-tree-nurseries/at\\_download/file](https://rngr.net/publications/proceedings/1979/soil-and-plant-analytical-services-for-tree-nurseries/at_download/file)
- South DB (2018) Fertilizer trials for bareroot nurseries in North America. *Reforesta* 5: 54-76. <https://dx.doi.org/10.21750/REFOr.5.07.53>
- South DB, Davey CB (1983) The southern forest nursery soil testing program. R8-TP-4. USDA Forest Service, Southern Region, Atlanta, GA: 140-170. [https://rngr.net/publications/1982-southern-nursery-conferences/the-southern-forest-nursery-soil-testing-program/at\\_download/file](https://rngr.net/publications/1982-southern-nursery-conferences/the-southern-forest-nursery-soil-testing-program/at_download/file)
- South DB, Zwolinski JB (1996) Chemicals used in southern forest nurseries. *South J Appl Forest* 20(3): 127-135. <https://doi.org/10.1093/sjaf/20.3.127>

- South DB (2021) Use of boron in conifer nurseries. *Reforesta* 12:56-97. <https://journal.reforestationchallenges.org/index.php/REFOR/article/download/141/137>
- South DB, Mitchell RJ, Dixon RK, Vedder M (1988) New-ground syndrome: an ectomycorrhizal deficiency in pine nurseries. *South J Appl Forest* 12(4): 234-239. <https://doi.org/10.1093/sjaf/12.4.234>
- South DB, Nadel RL, Enebak SA, Bickerstaff G (2017) Sulfur and lime affect soil pH and nutrients in a sandy *Pinus taeda* nursery. *Reforesta* (4): 12-20. <https://doi.org/10.21750/REFOR.4.02.41>
- South DB, Funk J, Davis CM (2018) Spring fumigation using totally impermeable film may cause ectomycorrhizal deficiencies at sandy loblolly pine nurseries. *Tree Planters' Notes* 61(1): 45-56. [https://rngr.net/publications/tpn/61-1/spring-fumigation-using-totally-impermeable-film-may-cause-ectomycorrhizal-deficiencies-at-sandy-loblolly-pine-nurseries/at\\_download/file](https://rngr.net/publications/tpn/61-1/spring-fumigation-using-totally-impermeable-film-may-cause-ectomycorrhizal-deficiencies-at-sandy-loblolly-pine-nurseries/at_download/file)
- Spilsbury R (1949) Maintenance of soil fertility. Forest Tree Nursery Meeting, Seattle, WA: 31-35. [https://rngr.net/publications/proceedings/1949/pdf.2005-12-09.2578998442/at\\_download/file](https://rngr.net/publications/proceedings/1949/pdf.2005-12-09.2578998442/at_download/file)
- Starkey T, Enebak S (2012) Foliar nutrient survey of loblolly and longleaf pine seedlings. Research Report 12-02. Auburn University Southern Forest Nursery Management Cooperative, Auburn University, AL: 11 p.
- Steinbeck K (1962) Effects of nutrients on slash pine seedlings grown in different media. Athens, GA: University of Georgia. 67 p. M.S. thesis.
- Steven HM (1928) Nursery investigations. *Forestry* 70: 31-11. <https://www.forestresearch.gov.uk/documents/6472/FCBU011.pdf>
- Stewart JH, Hite BH (1903) Commercial fertilizers: report for 1903. West Virginia Agricultural and Forestry Experiment Station Bulletin 92. <https://doi.org/10.33915/agnic.92>
- Stoeckeler JH (1949) Correction of soil acidity in conifer nurseries. Technical note 319. USDA Forest Service, Lake States Forest Experiment Station, St. Paul, MN: 1 p.
- Stoeckeler JH, Jones GW (1957) Forest nursery practice in the Lake States. Agriculture Handbook 110. USDA Forest Service, Washington, DC: 124 p. <https://babel.hathitrust.org/cgi/pt?id=uiug.30112019250551&view=1up&seq=1&skin=2021>
- Stoeckeler JH, Arneman HF (1960) Fertilizers in forestry. *Adv Agron* 12: 127-195. [https://doi.org/10.1016/S0065-2113\(08\)60082-6](https://doi.org/10.1016/S0065-2113(08)60082-6)
- Stone EL (1953) Magnesium deficiency of some northeastern pines. *Soil Sci Soc Am J* 17(3): 297-300. <https://doi.org/10.2136/sssaj1953.03615995001700030029x>
- Sucoff EI (1961) Potassium, magnesium, and calcium deficiency symptoms of loblolly and Virginia Pine seedlings. Station Paper NE-164. USDA Forest Service, Northeastern Forest Experiment Station, Upper Darby, PA: 18 p. [https://www.nrs.fs.fed.us/pubs/sp/sp\\_ne164.pdf](https://www.nrs.fs.fed.us/pubs/sp/sp_ne164.pdf)
- Sucoff EI (1962) Potassium, magnesium, and calcium requirements of Virginia pine. Station Paper NE-169. USDA Forest Service, Northeastern Forest Experiment Station, Upper Darby, PA: 16 p. <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.887.3525&rep=rep1&type=pdf>
- Sudworth GB (1900) The forest nursery: collection of tree seeds and propagation of seedlings. Bulletin 29. USDA, Division of Forestry. Washington, DC: 63 p. <https://ir.library.oregonstate.edu/downloads/0z708x74d>
- Sumner ME, Shahandeh H, Bouton J, Hammel J (1986) Amelioration of an acid soil profile through deep liming and surface application of gypsum. *Soil Sci Soc Am J* 50(5): 1254-1258. <https://doi.org/10.2136/sssaj1986.03615995005000050034x>
- Sung SS, Black CC, Kormanik TL, Zarnoch SJ, Kormanik PP, Counce PA (1997) Fall nitrogen fertilization and the biology of *Pinus taeda* seedling development. *Can J Forest Res* 27(9): 1406-1412. <https://doi.org/10.1139/x97-112>
- Swan HSD (1970) Relationships between nutrient supply, growth and nutrient concentrations in the foliage of black spruce and jack pine. Woodlands Reports 19. Pulp and paper Research institute of Canada. Pointe Claire. 46 p.
- Thomas GW, Jackson RM (1983) Growth responses of Sitka spruce seedlings to mycorrhizal inoculation. *New Phytologist* 95(2): 223-229. <https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1469-8137.1983.tb03488.x>
- Tillotson CR (1917) Nursery practice on the national forests. Bulletin 479. USDA. Washington, DC: 86 p. <https://babel.hathitrust.org/cgi/pt?id=uiug.30112019242087&view=1up&seq=1&skin=2021>

- Toumey JW (1916) Seeding and planting: a manual for the guidance of forestry students, foresters, nurserymen, forest owners, and farmers. John Wiley & Sons, NY.  
<https://babel.hathitrust.org/cgi/pt?id=uc2.ark:/13960/t12n50905&view=1up&seq=7&skin=2021>
- van den Driessche R (1963) Nursery experiments with Douglas fir. *The Commonwealth Forestry Review* 42(3): 242-254. <https://www.jstor.org/stable/42603050>
- van den Driessche R (1984) Soil fertility in forest nurseries. In: Duryea ML, Landis TD (eds) *Forest Nursery Manual*. MartinusNijhoff/Junk Publishers, The Hague, The Netherlands: 63-74.  
<https://rngr.net/publications/nursery-manuals/fnm/Chapter%207/>
- van den Driessche R (1989) Nutrient deficiency symptoms in container-grown Douglas-fir and white spruce seedlings. FRDA Report 100. Victoria, BC: B.C. Ministry of Forests. 29 p  
<https://www.for.gov.bc.ca/hfd/pubs/docs/frr/Frr100.htm>
- van den Driessche R (1991) Effects of nutrients on stock performance in the forest. In: van den Driessche (ed) *Mineral Nutrition of Conifer Seedlings*, CRC Press, Boca Raton: 229-260.
- VanderSchaaf C, McNabb K (2004) Winter nitrogen fertilization of loblolly pine seedlings. *Plant Soil* 265(1): 295-299. <https://doi.org/10.1007/s11104-005-0510-x>
- Van Goor CP (1963) Bemestingsvoorschriftvoornaaldhoutculturen. *Ned Bosh Tijdschr*35(3): 129-142.  
<https://edepot.wur.nl/417344>
- Van Goor CP (1970) Fertilization of conifer plantations. *Irish Forestry* 27(2): 68-80.  
<https://journal.societyofirishforesters.ie/index.php/forestry/article/download/9179/8322>
- van Lear DH, Smith WH (1972) Relationships between macro-and micronutrient nutrition of slash pine on three coastal plain soils. *Plant Soil* 36(1-3): 331-347. <https://doi.org/10.1007/BF01373488>
- Van Rees KCJ, Comerford NB, McFee WW (1990) Modeling potassium uptake by slash pine seedlings from low-potassium-supplying soils. *Soil Sci Soc Am J* 54(5): 1413-1421.  
<https://doi.org/10.2136/sssaj1990.03615995005400050034x>
- van Schöll L, Kuyper TW, Smits MM, Landeweert R, Hoffland E, Van Breemen N (2008) Rock-eating mycorrhizas: their role in plant nutrition and biogeochemical cycles. *Plant Soil* 303(1): 35-47.  
<https://core.ac.uk/download/pdf/29259822.pdf>
- Voigt GK (1955) The effect of applied fungicides, herbicides, and insecticides on the content of nutrient elements in tissue of coniferous seedlings. *Soil Sci Soc Am J* 19(2): 237-239.  
<https://doi.org/10.2136/sssaj1955.03615995001900020033x>
- Voigt GK, Stoeckeler JH, Wilde SA (1958) Response of coniferous seedlings to soil applications of calcium and magnesium fertilizers. *Soil Sci Soc Am J* 22(4): 343-345. <https://access.onlinelibrary-wiley-com.spot.lib.auburn.edu/doi/abs/10.2136/sssaj1958.03615995002200040022x>
- Wahlenberg WG (1930) Experiments in use of fertilizers in growing forest planting material at Savenac Nursery. Circular 125. USDA, Washington, DC: 38 p.
- Walker RF, McLaughlin SB (1997) Effects of acidic precipitation and ectomycorrhizal inoculation on growth, mineral nutrition, and xylem water potential of juvenile loblolly pine and white oak. *J Sustain Forest* 5(3-4): 27-49. [https://doi.org/10.1300/J091v05n03\\_03](https://doi.org/10.1300/J091v05n03_03)
- Wall RE (1978) Conifer seedling growth in limed peat in copper naphthenate-treated flats. *Tree Planters' Notes* 29(1): 18-21. [https://rngr.net/publications/tpn/29-1/29\\_1\\_18\\_21.pdf/at\\_download/file](https://rngr.net/publications/tpn/29-1/29_1_18_21.pdf/at_download/file)
- Wall MM (1994) Influence of fertilization on nutrient status and size of bare-root *Pinus taeda* L. seedlings. MS thesis, Texas A&M University, College Station. 98 p.  
<https://hdl.handle.net/1969.1/ETD-TAMU-1994-THESIS-W1875>
- Wang X, Zabowski D (1998) Nutrient composition of Douglas-fir rhizosphere and bulk soil solutions. *Plant Soil* 200(1): 13-20. <https://doi.org/10.1023/A:1004240315308>
- White EH, Comerford NB, Bickelhaupt DH (1980) Interpretation of nursery soil and seedling analysis to benefit nursery soil management. In: *Proceedings of the North American Forest Tree Nursery Soils Workshop*. Syracuse, NY: 269-287.  
<https://babel.hathitrust.org/cgi/pt?id=pst.000007190872&view=1up&seq=9&skin=2021>
- Wilde SA (1938) Soil fertility standards for growing northern conifers in forest nurseries. *J Agric Res* 57: 945-952. <https://naldc.nal.usda.gov/download/IND43969114/PDF>

- Wilde SA (1946) Forests soils and forests growth. Chronica Botanica Company, Waltham.  
<https://reader.library.cornell.edu/docviewer/digital?id=chla5566828>
- Wilde SA (1954) Reaction of soils; facts and fallacies. Ecology 35(1): 89-91.  
<https://doi.org/10.2307/1931409>
- Wilde SA (1958) Forest soils, their properties and relation to silviculture. Ronald Press, NY.  
<https://doi.org/10.2134/agronj1959.00021962005100100030x>
- Wilde SA, Kopitke JC (1940) Base exchange properties of nursery soils and the application of potash fertilizers. J Forest 38(4): 330-332. <https://doi.org/10.1093/jof/38.4.330>
- Wilde SA, Krause H (1959) Soil analysis in service of nursery practice. Tree Planters' Notes 37: 1-3.  
[https://rng.net/publications/tpn/10-3/PDF.2004-08-06.1516/at\\_download/file](https://rng.net/publications/tpn/10-3/PDF.2004-08-06.1516/at_download/file)
- Will GM (1961) Magnesium deficiency in pine seedlings growing in pumice soil nurseries. New Zealand Journal of Agricultural Research 4(1-2): 151-160.  
<https://www.tandfonline.com/doi/pdf/10.1080/00288233.1961.10419929>
- Will GM (1962) The uptake of nutrients from sterilised forest-nursery soils. New Zealand Journal of Agricultural Research 5(5-6): 425-432.  
<https://www.tandfonline.com/doi/pdf/10.1080/00288233.1962.10419939>
- Will GM (1985) Nutrient deficiencies and fertilizer use in New Zealand exotic forests. NZ For Res Inst Bull 97: 53 p.  
<https://scion.contentdm.oclc.org/digital/api/collection/p20044coll6/id/264/download>
- Will GM, Knight PJ (1968) Pumice soils as a medium for tree growth: pot trial evaluation of nutrient supply. New Zealand Journal of Forestry 13(1): 50-65.  
[https://nzjf.org.nz/free\\_issues/NZJF13\\_1\\_1968/C3FD6A3B-07E4-4B10-81C0-C8214E135DD2.pdf](https://nzjf.org.nz/free_issues/NZJF13_1_1968/C3FD6A3B-07E4-4B10-81C0-C8214E135DD2.pdf)
- Woodwell GM (1958) Factors controlling growth of pond pine seedlings in organic soils of the Carolinas. Ecol Monogr 28(3): 220-236. <https://doi.org/10.2307/1942241>
- Youngberg CT (1952) Maintenance of nursery soil fertility. In: Nursery practice committee. New Westminster, BC: 16-31.  
<https://nnp.rng.net/publications/proceedings/1952/maintenance-of-nursery-soilfertility/?searchterm=youngberg>
- Youngberg CT (1958) The uptake of nutrients by western conifers in forest nurseries. J Forest 56(5): 337-340. <https://doi.org/10.1093/jof/56.5.337>
- Youngberg CT (1984) Soil tissue analysis: tools for maintaining soil fertility. In: Duryea ML, Landis TD (eds) Forest nursery manual. MartinusNijhoff/Junk Publishers. The Hague, Netherlands, pp. 75-80. [https://link.springer.com/content/pdf/10.1007%2F978-94-009-6110-4\\_7.pdf](https://link.springer.com/content/pdf/10.1007%2F978-94-009-6110-4_7.pdf)
- Zarger TC (1964) Comparison of slowly and rapidly available nitrogen fertilizers for nursery production of pine seedlings. Tree Planters' Notes 66: 8-10.  
[https://rng.net/publications/tpn/15-3/comparison-of-slowly-and-rapidly-available-nitrogen-fertilizers-for-nursery-production-of-pine-seedlings/at\\_download/file](https://rng.net/publications/tpn/15-3/comparison-of-slowly-and-rapidly-available-nitrogen-fertilizers-for-nursery-production-of-pine-seedlings/at_download/file)
- Zhang J, George E (2010) Effect of the ectomycorrhizal fungus *Paxillus involutus* on growth and cation (potassium, calcium, and magnesium) nutrition of *Pinus sylvestris* L. in semi-hydroponic culture. J Plant Nutr 33(5): 736-751. <https://doi.org/10.1080/01904160903575956>